

**MINISTRY OF EDUCATION AND SCIENCE OF UKRAINE  
UZHHOROD NATIONAL UNIVERSITY  
FACULTY OF MEDICINE  
DEPARTMENT OF BIOCHEMISTRY AND PHARMACOLOGY**

**FUNCTIONAL BIOCHEMISTRY OF  
BLOOD, LIVER AND KIDNEYS**

**Manual for medical students**

Uzhhorod – 2021

**УДК 577.1(075.8)=111**

**Упорядники:**

- зав. кафедри біохімії та фармакології медичного факультету ДВНЗ «УжНУ», к.мед.н., доц. **Ростока Лариса Михайлівна,**
- асистент кафедри біохімії та фармакології медичного факультету ДВНЗ «УжНУ» **Сіткар Андрій Дмитрович**
- асистент кафедри біохімії та фармакології медичного факультету ДВНЗ «УжНУ» **Бурмістрова Яна Юріївна**

Методичний посібник для студентів складено у відповідності з вимогами освітньо-професійної програми підготовки магістра.

**РЕЦЕНЗЕНТИ:**

**Добрянська Е.С.** – к.мед.н., доцент кафедри фундаментальних медичних дисциплін медичного факультету №2 ДВНЗ «УжНУ»

**Дербак М.А.** – д.мед.н., професор кафедри пропедевтики внутрішніх хвороб медичного факультету ДВНЗ «УжНУ»

Затверджено на засіданні Вченої ради медичного факультету ДВНЗ «УжНУ»  
від 20.05.2021 р., протокол №7.

## **FOREWORD**

This manual provides information about functional biochemistry of blood, liver and kidneys in normal and some pathology conditions. The following topics are described: physical and chemical properties and composition of blood, respiratory function of erythrocytes, iron metabolism, function of blood plasma proteins, acid-base balance and its disorders, porphyrin metabolism, some questions enzyme diagnostics, integral role of liver in metabolism, bilirubin metabolism and biochemistry of jaundices, biotransformation of xenobiotics and endogenous metabolites, kidney functions, properties and composition of urine, water-mineral metabolism and its disorders, etc.

At the end of each chapter there are control questions and a list of information sources.

Perfect knowledge of the above topics is important for future doctors, because these chapters are directly related to clinical medicine and medical future practice.

# CHAPTER 1. BIOCHEMISTRY AND PATHOBIOCHEMISTRY OF BLOOD


## **Blood as a specialized tissue of the body, its composition. Functions of blood.**

**Blood** is fluid tissue composed of cells (formed elements of the blood) and an extracellular liquid medium. The overlying liquid (supernatant) of blood sample obtained on precipitation of the blood cells in the presence of an anticoagulant is called **blood plasma**. The plasma is an opalescent liquid containing all extracellular components of the blood. The blood cells account for about 45%, and the plasma, for about 55% of the blood volume. The clear liquid that separates from the blood when it is allowed to clot completely is called **blood serum**. Actually, the blood serum is the plasma from which fibrinogen has been removed in the process of clotting.

Suspended in the watery plasma are seven types of cells and cell fragments:

- 1) Red blood cells (RBCs) or erythrocytes;
- 2) Platelets or thrombocytes;
- 3) Five kinds of white blood cells (WBCs) or leukocytes: three kinds of granulocytes (neutrophils, eosinophils, basophils); two kinds of leukocytes without granules in their cytoplasm (lymphocytes, monocytes).

If one takes a sample of blood, treats it with an agent to prevent clotting, and spins it in a centrifuge, the red cells settle to the bottom the white cells settle on top of them forming the "buffy coat". The fraction occupied by the red cells is called the **hematocrit**. Normally it is approximately 45%. Values much lower than this are a sign of anemia.



**Figure 1.** Separation of blood into its basic components (Janson, 2012)

### **Major functions of blood:**

- 1) Respiration (transport of O<sub>2</sub> from the lungs to the tissues and CO<sub>2</sub> from tissues to lungs).
- 2) Transport function (transport of different substances).
- 3) Excretion (transport of metabolic products to the kidneys, lungs, skin and intestine for removal).
- 4) Maintenance of the normal acid-base balance in the body.

- 5) Regulation of water balance through the effects of blood on the exchange of water between the circulating fluid and the tissue fluid.
- 6) Regulation of body temperature by means of distribution of body heat.
- 7) Defense against infection by the white blood cells and circulating antibodies.
- 8) Transport of hormones and regulation of metabolism.

**Blood as source for medicinal preparations.** The blood is used as a raw material for producing a variety of medicinal preparations which, by their therapeutic applications, are divided into four groups:

- 1) systemic effect agents (albumin, protein, native blood plasma);
- 2) immunologically active preparations (gamma-globulin, antistaphylococcic, interferon)
- 3) hemostatic preparations (antihemophilic plasma, thrombin, fibrin sponge, fibrin film, fibrinogen);
- 4) antianemic and stimulating preparations (polyobolin – dry powdered protein components of blood plasma, eryheme-dehydrated hemolyzate of erythrocytes, etc.).

### **Physical and chemical properties of blood. Inorganic components of blood.**

Normally the average blood volume is 5200 ml in men and 3900 ml in women. The blood plasma accounts for about 55% of the total volume. The erythrocytes constitute a major fraction of blood cells and account for 44% of the total blood volume. Other blood cells account only 1%. The relative **density** of whole blood is 1,05-1,064, of blood plasma – 1,024-1,030, of blood cells – 1,08-1,097. **Viscosity** of blood is 4-5 fold that of water. It is provided by means of high content of proteins and erythrocytes.

An essential physico-chemical characteristic of the blood is the **osmotic pressure** of blood plasma. It is provided by osmotic concentration that is by the sum total of all the blood particles per unit of volume. At a body temperature of 37°C, the blood plasma osmotic pressure is about 7,6 atm. (768-818 kPa). The major contributors to this value are NaCl and other low-molecular weight substances, contained in the blood. Osmotic pressure constancy provides the normal transport of substances from blood to tissues and back, promotes the stability of erythrocytes. The part of osmotic pressure, which is provided by proteins, is called **oncotic (or colloid osmotic) pressure**. Oncotic pressure accounts for about 0,03 atm. (0,5% of osmotic pressure). But oncotic pressure is very important, as proteins cannot penetrate through semipermeable membrane and therefore oncotic pressure facilitates the reverse stream of fluid to venous part of capillaries.

#### **Mineral substances of blood:**

**Sodium** is a major osmotically active ion in the extracellular space. In **hypernatremia**, a syndrome associated with the organism's hyperhydration is commonly observed to develop. The accumulation of excessive sodium in the blood plasma occurs in a specific renal disease known as parenchymatous nephritis, in patients with congenital cardiac insufficiency, also in primary (or true) and secondary hyperaldosteronism. **Hyponatremia** is accompanied by dehydration of the organism.

**Potassium.** The potassium level in the cells is much higher as compared to the extracellular space. **Hyperkalemia** is observed in acute renal insufficiency or in hypofunction of adrenal cortex. By contrast, an increased production of aldosterone by adrenal cortex leads to **hypokalemia**. The progressive hypokalemia leads to grave disturbances of cardiac performance. Occasionally, a decreased level of potassium in the

blood serum was observed as a side effect on administration of large therapeutic doses of adrenal cortex hormones to patients.


**Calcium.** In tumoral lesions of bone tissue, hyperplasia, or parathyroid adenoma, a marked **increase of calcium** level in the blood plasma is observed. The state of **hypocalcemia** is observed in hypoparathyroidism. The hypofunction of parathyroid gland results in a drastic drop of calcium concentration in the blood, with the eventual development into a convulsive state (tetany). Hypocalcemia is also observed in rickets, sprue, obstructive jaundice, nephroses, glomerulonephritis.

<b>Chemical composition of blood plasma</b>	
<p><b>I. Proteins</b></p> <ol style="list-style-type: none"> <li>1. Total protein 65-85 g/L</li> <li>2. Albumins 35-50 g/L</li> <li>3. Globulins 25-35 g/L</li> <li>4. Fibrinogen 2.0-7.0 g/L</li> <li>5. Haptoglobin 0.28-1.90 g/L</li> <li>6. Prothrombin 10-15 mg/dL</li> <li>7. Plasminogen 1.4-2.8 <math>\mu\text{mol/L}</math> (20-40 mg/dL)</li> <li>8. Transferrin 19.3-45.4 <math>\mu\text{mol/L}</math> (170-400 mg/dL)</li> <li>9. Ceruloplasmin 1.52-3.31 <math>\mu\text{mol/L}</math> (23-50 mg/dL)</li> <li>10. <math>\beta</math>-Lipoproteins 3.0-6.0 g/L (300-600 mg/dL)</li> </ol> <p>HDL – high density lipoprotein (<math>\alpha</math>-LP) 1.063-1.210 mmol/L (80-400 mg/dL)</p> <p>LDL – low density lipoprotein (<math>\beta</math>-LP) 1.006-1.063 mmol/L (360-640 mg/dL)</p>	<p><b>II. Enzymes</b></p> <ol style="list-style-type: none"> <li>1. Alanine aminotransferase (ALT) 0.16-0.68 mmol/h·L or (15-75 IU/L) (glutamate pyruvate transferase, GPT)</li> <li>2. Aspartate aminotransferase (AST) 0.10-0.45 mmol/h·L or (10-50 IU/L) (glutamate oxaloacetate transferase, GOT)</li> <li>3. Lactate dehydrogenase 0.8-4.0 mmol/h·L</li> <li>4. Creatine kinase &lt; 1.2 mmol/h·L or (&lt; 90 IU/L)</li> <li>5. Fructose-biphosphate aldolase (F-1,6-PA) 3.6-21.8 mmol/h·L</li> <li>6. Acetylcholine esterase 160-340 mmol/h·L</li> <li>7. <math>\alpha</math>-Amylase 15-30 g/h·L or (&lt; 300 IU/L)</li> <li>8. Alkaline phosphatase 30-150 IU/L</li> <li>9. Acidic phosphatase &lt; 62 nkat/L</li> <li>10. <math>\gamma</math>-Glutamyl transferase (<math>\gamma</math>GT or GGT) &lt; 60 IU/L</li> </ol>
<p><b>III. Nonproteinic nitrogenous compounds</b></p> <ol style="list-style-type: none"> <li>1. Nitrogen residual (nonproteinic) 19.5-30.0 mmol/L</li> <li>2. Nitrogen of amino acids 3.5-5.5 mmol/L</li> <li>3. Creatine 15-70 mmol/L</li> <li>4. Creatinine 60-150 <math>\mu\text{mol/L}</math></li> <li>5. Urea 3.3-6.7 mmol/L</li> <li>6. Uric acid 0.1-0.4 mmol/L</li> <li>7. Bilirubin total 8-20 <math>\mu\text{mol/L}</math></li> <li>8. N-Acetylneuraminic acid 1.8-2.2 mmol/L</li> <li>9. Histamine 17.99-71.94 nmol/L (0.2-0.8 ng/dL)</li> <li>10. Adrenalin 1.91-2.46 nmol/L (0.35-0.45 ng/L)</li> <li>11. Serotonine 0.3-1.7 <math>\mu\text{mol/L}</math> (5.0-30.0 <math>\mu\text{g/dL}</math>)</li> <li>12. Thyroxine 64.36-141.59 nmol/L (5-11 ng/dL)</li> </ol>	<p><b>IV. Carbohydrates and metabolites</b></p> <ol style="list-style-type: none"> <li>1. Glucose 2.8-6.0 mmol/L</li> <li>2. Lactate 0.5-2.0 mmol/L</li> <li>3. Pyruvate &lt; 0.1 mmol/L</li> <li>4. Citric acid 88.5-156.1 <math>\mu\text{mol/L}</math> (1.7-3.0 mg/dL)</li> </ol> <p><b>V. Lipids and metabolites</b></p> <ol style="list-style-type: none"> <li>1. Total lipids 4.0-8.0 g/L</li> <li>2. Triacylglycerides 0.5-2.1 mmol/L</li> <li>3. Total phospholipids 2.0-3.5 mmol/L</li> <li>4. Total cholesterol 4.0-8.6 mmol/L</li> <li>5. Free fatty acids 0.3-0.8 mmol/L</li> <li>6. Ketone bodies 100-600 <math>\mu\text{mol/L}</math></li> </ol>
<p><b>VI. Mineral components</b></p> <ol style="list-style-type: none"> <li>1. Sodium (<math>\text{Na}^+</math>) 135-155 mmol/L</li> <li>2. Potassium (<math>\text{K}^+</math>) 3.6-5.0 mmol/L</li> <li>3. Chlorides (<math>\text{Cl}^-</math>) 97-108 mmol/L</li> <li>4. Calcium (<math>\text{Ca}^{2+}</math>) 2.25-2.75 mmol/L</li> <li>5. Phosphate inorganic 0.8-1.4 mmol/L</li> <li>6. Magnesium (<math>\text{Mg}^{2+}</math>) 0.7-1.0 mmol/L</li> <li>7. Sulphates 0.4-0.6 mmol/L</li> <li>8. Iron (Fe) 14-32 <math>\mu\text{mol/L}</math> (65-175 <math>\mu\text{g/dL}</math>)</li> <li>9. Copper (Cu) 12-19 <math>\mu\text{mol/L}</math></li> <li>10. Zinc (Zn) 12-20 <math>\mu\text{mol/L}</math></li> <li>11. Ammonia 10-47 <math>\mu\text{mol/L}</math></li> </ol>	<p><b>Indices of blood</b></p> <ol style="list-style-type: none"> <li>1. Hemoglobin: males 130-180 g/L (13-18 g/dL), females 120-160 g/L (12-16 g/dL)</li> <li>2. Hydrogen ion: arterial blood 35-46 nmol/L (pH=7.36-7.44) (38°C)</li> <li>3. Oxygen (<math>\text{pO}_2</math>) in arterial blood: 11-15 kPa (85-105 mm Hg)</li> <li>4. Bicarbonate total (<math>\text{CO}_2</math>) 22-30 mmol/L</li> <li>5. Carbon dioxide (<math>\text{pCO}_2</math>) in arterial blood: 4.5-6.0 kPa (35-46 mm Hg)</li> </ol>

## Buffer blood systems. Acid-base balance, its regulation. Acidosis and alkalosis.

**Acid-base balance** is the relation between concentrations of hydrogen and hydroxyl ions in liquids of organism. This balance is characterized by the hydrogen ion concentration (in nmoles per liter), or by the pH value which is the negative logarithm (to base 10) of hydrogen ion concentration. Blood hydrogen ion concentration  $[H^+]$  is maintained with tight limits in health. Normal levels lie between 35-45 nmol/L (pH 7,35-7,45). Values greater than 120 nmol/l (pH 6,92) or less than 20 nmol/L (pH 7,7) are usually incompatible with life. The total amount of hydrogen ion produced each day in this way is of the order of 60 mmoles. If all of this was be diluted in the extracellular fluid (~14 liters),  $[H^+]$  would be 4 mmol/L or 100 000 times more acid than normal. This just does not happen, as all the hydrogen ions produced are efficiently excreted in urine.

The „first line” of defense from changing pH is buffer systems. The buffer system is a conjugated acid-base pair composed of a donor and an acceptor of hydrogen ions (protons). The acid-base balance of a buffer solution is described by the Henderson-Hasselbach equation:  $pH = pK_a + \log [\text{proton acceptor}]/[\text{proton donor}]$ . The major buffer systems of blood are bicarbonate, phosphate, protein, and especially hemoglobin systems.



**Figure 2.** Buffer systems in the plasma (Koolman, 2005)

**The bicarbonate buffer system** is powerful and perhaps the most controllable system of both the extracellular fluid and blood. Bicarbonate buffer system accounts for about 10% of the total buffering capacity of blood.



$$[HCO_3^-] = 27 \text{ mmol/L}$$

$$[H_2CO_3] = 1,35 \text{ mmol/L}$$

$$[HCO_3^-] / [H_2CO_3] = 20/1$$


Bicarbonate buffer system functions as an effective medium regulator within a close range of pH 7,4. The limit to the effectiveness of the bicarbonate system is the initial concentration of bicarbonate. Only when all the bicarbonate is used up the system has no further buffering capacity. The acid-base status of patients is assessed by consideration of the bicarbonate system of plasma. The extracellular fluid contains a large amount of bicarbonate about 24 mmol/L.

**Phosphate buffer systems** is a conjugated acid-base pair composed of ion  $\text{H}_2\text{PO}_4^-$  (proton donor) and ion  $\text{HPO}_4^{2-}$  (proton acceptor).

$$[\text{HPO}_4^{2-}] / [\text{H}_2\text{PO}_4^-] = 4/1$$

Phosphate buffer system accounts about 1% of the blood buffering capacity. Nonetheless, in the tissues, especially in kidneys, this system is a major one. This buffer is operative within a range of pH variation from 6,1 to 7,7.

**Protein buffer system** is of minor importance for maintaining the acid-base balance in the blood plasma as compared to other buffer systems. Proteins form a buffer system owing to the occurrence of acid and basic groups in their molecules:



The protein buffer system of blood plasma is effective within a pH range 7,2-7,4.

**Hemoglobin buffer system** is the most powerful buffer system of blood. This buffer system accounts for about 75% of the total buffering capacity of blood. The involvement of hemoglobin in the control of blood pH is primarily associated with the function of hemoglobin in the transport of oxygen and carbon dioxide. The dissociation constancy of acidic hemoglobin groups is liable to variation depending on the degree of hemoglobin saturation with oxygen. Hemoglobin, on its uptake of oxygen, becomes a stronger acid ( $\text{H}^+ + \text{HbO}_2$ ). By contrast hemoglobin without oxygen is a very weak organic acid (HHb).

**Disturbances of acid-base balance.**

**Acidosis:** the hydrogen ion concentration in the blood is above the normal; level of pH below 6,92 ( $\uparrow[\text{H}^+] 120 \text{ nmol/l}$ ) causes death.

**Alkalosis:** the hydrogen ion concentration in the blood is lower the normal; level of pH above 7,7 ( $\downarrow[\text{H}^+] 20 \text{ nmol/l}$ ) is incompatible with life.

Blood gas results			
Metabolic acid-base disorders		Respiratory acid-base disorders	
<b><math>[\text{H}^+]</math> elevated</b>	<b><math>[\text{H}^+]</math> decreased</b>	<b><math>\text{H}^+</math> elevated</b>	<b><math>\text{H}^+</math> decreased</b>
↓	↓	↓	↓
<b>Acidosis</b>	<b>Alkalosis</b>	<b>Acidosis</b>	<b>Alkalosis</b>
↓	↓	↓	↓
<b><math>\text{HCO}_3^- \downarrow</math></b>	<b><math>\text{HCO}_3^- \uparrow</math></b>	<b><math>\text{pCO}_2 \uparrow</math></b>	<b><math>\text{pCO}_2 \downarrow</math></b>
↓	↓	↓	↓
<b>metabolic acidosis</b>	<b>metabolic alkalosis</b>	<b>respiratory acidosis</b>	<b>respiratory alkalosis</b>

**Metabolic acidosis** is the most common form of disturbed acid-base balance. It is associated with the accumulation of organic acids in the tissues and the blood. Causes of metabolic acidosis:

- 1) Renal disease (acute and chronic glomerulonephritis; acute and chronic pyelonephritis). The decrease of excretion of protons by kidney.
- 2) Ketoacidosis (imperfect oxidation of lipids: in diabetes mellitus, starvation, fever; ketogenic diet).
- 3) Lactate acidosis (imperfect oxidation of carbohydrates, lung diseases, cardiovascular diseases, different types of hypoxia).
- 4) Certain causes of over dosage of poisoning (salicylate  $\rightarrow$  lactate; methanol  $\rightarrow$  formate; ethylene glycol  $\rightarrow$  oxalate).



5) Chronic diarrhea or intestinal fistula.

**Clinical effects of metabolic acidosis.** The compensatory response to metabolic acidosis is hyperventilation since the increased  $[H^+]$  acts as a powerful stimulant of the respiratory centre. The deep, rapid and gasping respiratory pattern is known as Kussmaul breathing. Hyperventilation is the appropriate physiological response to acidosis and it occurs rapidly. A raised  $[H^+]$  leads to increased neuromuscular irritability. There is a hazard of arrhythmias progressing to cardiac arrest, and this is made more likely by the presence of hyperkalemia which will accompany the acidosis.

**Metabolic alkalosis** is due to loss of a large amount of acid equivalents (for example, in noncontrolable vomiting), the accumulation of base equivalents in tissues (for example in tetany), a wrong correction for metabolic acidosis, high doses of glucocorticoids. The condition may be due to:

- 1) Loss of hydrogen ion in gastric fluid during vomiting.
- 2) Ingestion of an absorbable alkali such as sodium bicarbonate.
- 3) Potassium deficiency.

In severe potassium depletion, often a consequence of diuretic therapy, hydrogen ions are retained inside cells to replace the missing potassium ions. In the renal tubule more hydrogen ions, rather than potassium, are exchanged for reabsorbed sodium. So, despite there being an alkalosis, the patient passes an acid urine. This is often referred to as a "paradoxical" acid urine, because in other causes of metabolic alkalosis urinary  $[H^+]$  usually falls.

**Clinical effects of metabolic alkalosis.** The clinical effects of alkalosis include hypoventilation, confusion and eventually coma. Muscle cramps, tetany and parasthesia may be consequence of a decrease in the unbound plasma calcium concentration which is a consequence of the alkalosis.

Recognizing primary **metabolic acid-base disorders** by inspecting the  $HCO_3^-$  concentration.

**Respiratory acidosis** develops because of a reduced minute respiratory volume (for example, in bronchiole asthma, edema of lungs, emphysema, atelectasis and traumatic asphyxia). Respiratory acidosis may be acute or chronic. Examples of acute, and uncompensated, respiratory acidosis are:

- 1) choking;
- 2) bronchopneumonia;
- 3) acute exacerbation of asthma.

Chronic respiratory acidosis is usually a long-standing condition and is accompanied by maximal renal compensation. Examples of chronic respiratory disorders are:

- 1) chronic bronchitis;
- 2) emphysema.

**Respiratory alkalosis** arises from a sharply intensified pulmonary ventilation (in inhalation of pure oxygen, in compensatory dyspnea). Respiratory alkalosis is much less common than acidosis. Examples are:

- 1) hysterical over breathing;
- 2) mechanical over ventilation;
- 3) raised intracranial pressure or hypoxia, both of which may stimulate the respiratory centre.

**Blood plasma proteins. Pathoproteinemia. Acute phase proteins.  
Lipoproteins. Non-protein organic compounds of blood. Azotemia.**

Proteins account for 6,5-8,5% out of the total 9-10% of dry blood plasma residue.

Total proteins – 65-85 g/L

Albumins – 40-50 g/L    Globulins – 20-40 g/L    Fibrinogen – 1,5-3,5 g/L

The most of blood serum proteins are synthesized in liver, but some of them are formed in other tissues. For example,  $\gamma$ -globulins are synthesized by lymphocytes; peptide hormones are mainly secreted by endocrine glands; peptide hormone erythropoietin is formed by kidney cells. Almost all the blood plasma proteins, with the exception of albumin, are glycoproteins.

Group	Protein	$M_r$ , in kDa	Function
Albumins:	Transthyretin Albumin: 45 g · l <sup>-1</sup>	50-66 67	Transport of thyroxin and triiodothyronin Maintenance of osmotic pressure; transport of fatty acids, bilirubin, bile acids, steroid hormones, pharmaceuticals and inorganic ions.
$\alpha_1$ -Globulins:	Antitrypsin Antichymotrypsin Lipoprotein (HDL) Prothrombin  Transcortin  Acid glycoprotein Thyroxin-binding globulin	51 58-68 200-400 72  51  44 54	Inhibition of trypsin and other proteases Inhibition of chymotrypsin Transport of lipids Coagulation factor II, thrombin precursor (3.4.21.5)  Transport of cortisol, corticosterone and progesterone Transport of progesterone Transport of iodothyronins
$\alpha_2$ -Globulins:	Ceruloplasmin Antithrombin III Haptoglobin Cholinesterase (3.1.1.8) Plasminogen  Macroglobulin  Retinol-binding protein Vitamin D-binding protein	135 58 100 ca. 350 90  725  21 52	Transport of copper ions Inhibition of blood clotting Binding of hemoglobin Cleavage of choline esters Precursor of plasmin (3.4.21.7), breakdown of blood clots  Binding of proteases, transport of zinc ions  Transport of vitamin A Transport of calciols
$\beta$ -Globulins:	Lipoprotein (LDL) Transferrin Fibrinogen Sex hormone-binding globulin Transcobalamin C-reactive protein	2.000-4.500 80 340  65 38 110	Transport of lipids Transport of iron ions Coagulation factor I  Transport of testosterone and estradiol Transport of vitamin B <sub>12</sub> Complement activation
$\gamma$ -Globulins:	IgG IgA IgM IgD IgE	150 162 900 172 196	Late antibodies Mucosa-protecting antibodies Early antibodies B-lymphocyte receptors Reagins

**Figure 3.** Plasma proteins (Koolman, 2005)

**Functions of blood proteins:** they take part in blood clotting; they provide viscous properties of blood; maintaining acid base balance; transport function; protective function; reserve of amino acids; regulation function; they maintain the oncotic pressure; maintaining a needed level of cations in blood.

**Albumin** level in blood plasma protein is 35-50 g/L. Albumins make up approximately 60% of the total plasma protein. Functions of albumins:

1) Albumins are responsible for 75 – 80% of oncotic pressure of human's plasma. The decreasing albumin concentration below 30 g/L leads to edema.

2) Transport function. It transports free fatty acids, calcium, certain steroids hormones, bilirubin, copper, different drugs etc.

**Globulins:** are divided into  $\alpha_1$ -globulins (3-6 g/L),  $\alpha_2$ -globulins (4-9 g/L),  $\beta$ -globulins (6-11 g/l) and  $\gamma$ -globulins (7-15 g/L). They perform transport and protective functions.

#### **$\alpha$ -Globulins:**

**Haptoglobin** (Hp) (is component of  $\alpha_2$ -globulin fraction). This glycoprotein binds extracorporeal hemoglobin. The haptoglobin-hemoglobin complex can be absorbed by the macrophage system and cannot pass the glomerulus of the kidney. Thus Hp prevents a loss of free hemoglobin by kidney and provides the conservation and reutilization of iron.

**Ceruloplasmin** ( $\alpha_2$ -globulin) is a blue, copper-containing (0,32%) glycoprotein found in mammalian blood plasma. It contains about 3% of total amount of copper in organism and more 90% Cu of blood plasma. Ceruloplasmin exhibits a weakly pronounced catalytic activity in the oxidation of ascorbic acid, adrenaline, dihydrophenylalanine and a number of other compounds, and ferroxidase activity ( $\text{Fe}^{2+} \rightarrow \text{Fe}^{3+}$ ). It is antioxidant. In Wilson's diseases the concentration of ceruloplasmin in the blood plasma is significantly lowered which a major diagnostic test for this pathology.

**$\alpha_1$ -Antitrypsin** can inhibit trypsin and other proteolytic enzymes. The level of trypsin inhibitors is increased in inflammatory processes, in pregnancy and in a number of other states of organism. In inflammatory process the level of  $\alpha_1$ -antitrypsin increases in result of stimulation of its synthesis in hepatocytes. In acute pancreatitis the enhanced level of  $\alpha_1$ -antitrypsin arises from delivery of active pancreatic proteinases. Deficiency of  $\alpha_1$ -antitrypsin is associated with emphysema and one type of liver diseases ( $\alpha_1$ -antitrypsin deficiency liver disease).

**$\alpha_2$ - Macroglobulin** is a large plasma glycoprotein (720 kDa). It is inhibitor of serine-, thiol-, carboxy- and metal proteinases. Therefore it is involved in regulation of blood clotting, immunologic processes, inflammatory processes. In addition, it binds many cytokines (eg. platelet-derived growth factor, transforming growth factor- $\beta$ , etc.) and is involved in targeting them toward tissues.

#### **$\beta$ -Globulins:**

**C-reactive protein** is able to form a precipitate with the somatic C polysaccharide of pneumococcus C-reactive protein does not occur in the blood serum of healthy organisms. It is detected in many pathologic states attendant to inflammation and necrosis of the tissues. This protein is „acute phase” protein.

**Hemopexin** binds free heme.

**Transferrin** is glycoprotein with a molecular mass of approximately 80 kDa. Transferrin plays a central role in the body's metabolism of iron, because it transports iron (2 moles of  $\text{Fe}^{3+}$  per moll of transferrin) in the circulation to sites where iron is required, e.g. from the gut to the bone marrow and other organs.


#### **$\gamma$ -Globulins:**

**Cryoglobulin** is absent in the blood serum of healthy individuals. It is found only in pathologic states. A specific feature of this protein is its solubility at a temperature 37°C and ability to form a precipitate or a gel in decreasing temperature to 4°C. It is detected in blood

serum in myeloma, nephrosis, cirrhosis of the liver, rheumatism, lymphosarcoma, leukosis and other diseases.

**Interferons** are specific proteins synthesized on the organism's cells invaded by virus. Interferon can inhibit viral multiplication in the cells however it has no effect on the viral particles that have been formed in the cell. Interferon is easy to leave the cell and to enter the blood stream in which it is carried over to tissues and organs. There are 3 types of interferons: IFN- $\alpha$ , IFN- $\beta$ , and IFN- $\gamma$ . IFN- $\alpha$  are mainly synthesized by leukocytes; IFN- $\beta$  by fibroblasts; IFN- $\gamma$  by T- and B-lymphocytes.


**Immunoglobulins** (humoral antibodies) are synthesized mainly in plasmocytes, specialized cells of B-cell lineage that synthesize and secrete immunoglobulins into plasma. All immunoglobulin molecules consist of 4 polypeptide chains, which are linked by disulfide bonds: two identical light (L) chains and two identical heavy (H) chains. L-chains have molecular mass 23000 Da. They are common to all the classes of immunoglobulins. They are two types: kappa ( $\kappa$ ) and lambda ( $\lambda$ ). A given immunoglobulin molecule contains or two identical  $\kappa$ , or two  $\lambda$  chains. Heavy chains have molecular mass 50000-75000 Da. Five types of heavy chains exist:  $\alpha$  (alpha),  $\gamma$  (gamma),  $\mu$  (mu),  $\delta$  (delta),  $\epsilon$  (epsilon). The type of H chain determines the class of immunoglobulin and thus its effector function. There are five immunoglobulin classes: IgG ( $\gamma$ -chains), IgA ( $\alpha$ -chains), IgM ( $\mu$ -chains), IgD ( $\delta$ -chains), IgE ( $\epsilon$ - chains). Every light and heavy chain consist of 2 segments: variable (V) and constant (C). The constant regions of immunoglobulin molecules are responsible for the class specific effectors functions of the different immunoglobulin molecules, e.g. complement fixation or placental transfer.



**Figure 4.** Classes of immunoglobulins (Koolman, 2005)

IgGs (7-20 g/L) and IgMs (0,5-2 g/L) are basic classes of immunoglobins. They realize humoral immune response on the incorporation of foreign antigens. IgMs participate in the primary immune response, they activate complement system. IgGs participate in the secondary immune response, activate a complement system. IgG is the only immunoglobulin which is able to pass placenta. IgA (0,7-1,5 g/L) is antibody of other biological liquids and secretions (secretions of mucous, lungs). IgD (0,000001-0,0003 g/L) and IgE (0,02-0,02g/L) are minor components of blood serum. IgE participate in allergic reactions.

In clinical practice, there have been reported states characterized by alteration in both the total content of blood plasma proteins and the percentage of individual protein fractions.



**Figure 5.** Electrophoresis (Koolman, 2005)

**Hypoproteinemia** (a decrease in the total concentration of blood plasma proteins) is usually linked with the decreasing albumins. Hypoproteinemia occurs:

- 1) in nephrotic syndrome;
- 2) in liver disease (acute atrophy of the liver, toxic hepatitis, and other states);
- 3) in a drastically increased permeability of the capillary wall,
- 4) in protein deficiency (affected gastrointestinal tract, carcinoma, etc).

**Paraproteinemia** is the occurrence in the blood plasma of proteins, normally untypical to the healthy organism (for example, in myeloma). In the blood serum of patients with myeloma specific „myelomatous” proteins are detected.

**Hyperproteinemia** is a pathologic condition manifested by an increased content of blood plasma proteins. Hyperproteinemia:

- 1) relative (it is caused by loss of liquid by organism (diarrhea in children; vomiting, due to an obstruction of the upper small intestine, or by extensive burns);
- 2) absolute (it is caused by an elevated level of  $\gamma$ -globulins).

**Dysproteinemia** is the changing ratio of individual protein fractions, while the total protein content in the blood serum is normal.  $\gamma$ -Globulin fraction is increased in chronic inflammation, chronic polyarthritis etc.  $\alpha_2$ -Globulin fraction is increased in acute infections, acute rheumatism. The level of some proteins may be sharply raised in acute inflammatory processes and some other pathologic states (trauma, burns, myocardial infarction). Those proteins are called «**acute phase**» proteins, because they take part in development of inflammatory reaction of organism. Main inducer of the synthesis of the most acute phase proteins in hepatocytes is interleukin-1 liberated by mononuclear phagocytes. Haptoglobin, C-reactive protein,  $\alpha_1$ -antitrypsin, acid  $\alpha_1$ - glycoprotein, fibrinogen belong to proteins of acute phase.


## Blood plasma lipoproteins.

To transport through the blood water-insoluble lipids from one tissue to another special transport forms exist – lipoproteins. There are 5 classes of blood lipoproteins:

1. Chylomicrons;
2. VLDL (Very Low Density Lipoproteins);
3. IDL (Intermediate Density Lipoproteins);
4. LDL (Low Density Lipoproteins);
5. HDL (High Density Lipoproteins).


Despite their differences in lipid and protein composition, all lipoproteins share common structural features, notably they have a spherical shape, and consist of a core of triacylglycerols or/ and cholesterol esters surrounded by a single layer of phospholipids, into which a mixture of cholesterol and proteins (apoproteins) is inserted.

Note that the phospholipids and cholesterol are oriented with their polar head groups facing outward to interact with solvent water and thus shield the hydrophobic lipids inside from the water outside. The proteins also contribute to the formation of lipoprotein polar surface, but additionally act as cofactors for enzymes, which take part in lipoprotein metabolism, and function as recognition sites for the various lipoprotein receptors throughout the body.



**Figure 6.** Composition of lipoprotein complexes (Koolman, 2005)

**Chylomicrons** serve to transport triacylglycerols and cholesterol esters from the intestines to other tissues and eventually to the liver. Chylomicrons are synthesized in intestinal epithelial cells. They are the largest and least dense of the blood lipoproteins because they have the most triacylglycerol (about 90%) and rather low content of protein. Their triacylglycerols and cholesterol are derived from the dietary lipids, and their major protein is apo B-48. Chylomicrons enter the lymphatic system and travel through the lymph into the blood. Additional apoproteins (apo CII and apo E) are transferred to nascent chylomicrons from HDL, and mature chylomicrons are formed. In capillaries of peripheral tissues, particularly adipose and muscle, chylomicrons become the target for lipoprotein lipase. Lipoprotein lipase is activated by apo CII and hydrolyzes triacylglycerols to fatty acids and glycerol. Fatty acids are taken up and used by peripheral cells. As chylomicrons are degraded, chylomicron remnants are formed. Remnants interact with apo E receptors of hepatocytes, undergo endocytosis and lysosomal degradation.



**Figure 7.** Transport of lipoproteins (Koolman, 2005)

**VLDL**, **IDL** and **LDL** form a group of related particles that deliver endogenous triacylglycerols and cholesterol from the liver to the peripheral tissues. VLDL is synthesized in the liver. Its' core is formed from triacylglycerols (about 55%) and cholesterol esters. The phospholipids, cholesterol and apoprotein apo B-100 form the coat. VLDL particles are released into the blood and get additional apoproteins (apo CII and apo E) from HDL. In peripheral tissues, VLDL triacylglycerols are hydrolyzed by lipoprotein lipase, and VLDL is converted to IDL. Most of IDL return to the liver, bind to apo E receptors, undergo endocytosis and lysosomal degradation. About 25% of IDL in liver sinusoids are acted on by hepatic lipase. Further hydrolysis of triacylglycerols and removal of additional apoproteins results in LDL formation. LDLs react with apo B-100 receptors (LDLreceptors) on the cells

of various tissues, are taken up by endocytosis and decomposed in lysosomes. LDL is regarded as the vehicle delivering cholesterol to the peripheral tissues. Three regulatory mechanisms are used by cells to prevent excessive accumulation of cholesterol:

- 1) Cholesterol inhibits HMG-CoA reductase, the key enzyme of de novo cholesterol biosynthesis.
- 2) Cholesterol activates acyl-CoA:cholesterol acyltransferase (ACAT), which converts free cholesterol to cholesterol ester (cholesterol storage form).
- 3) Cholesterol inhibits synthesis of LDL-receptors thus reduces the amount of cholesterol taken up by the cell from the blood.

The removal of modified LDLs from the bloodstream occurs due to macrophages through their scavenger receptors. Scavenger receptors are not down-regulated by cholesterol, and unlimited uptake of LDL particles transforms macrophages to the “foam cells”. The accumulation and death of such foam cells in the intima of arteries cause atherosclerotic plaque formation, with cholesterol being its chief chemical constituent. Therefore, LDLs are referred to as “bad cholesterol”, and prolonged elevation of LDL levels followed by their oxidative and other modifications lead to atherosclerosis.

**HDL** is synthesized by the liver and released into the blood as disc-shaped particles. The major constituents of nascent HDL are phospholipids and proteins (apo A, C, D, E). HDLs participate in metabolism of chylomicrons and VLDLs providing apoproteins required for their metabolism. Besides, HDL takes cholesterol from the peripheral tissues and transports it back to the liver. Excessive cholesterol from the cells may be passed to HDL, converted to cholesterol ester by lecithin:cholesterol acyltransferase (LCAT) and shifted to the interior of the particle. LCAT is activated by apo A1. As cholesterol esters accumulate in the core of the lipoprotein, the particle becomes spherical (mature HDL). With the aid of apo D cholesterol esters partly may be transferred to the chylomicron remnants, IDL or other lipoproteins. Finally HDLs get unloaded of cholesterol in the liver, where it may be converted to the bile acids and secreted into the bile. Therefore, HDLs are often referred to as “good cholesterol” because they function to deliver cholesterol from peripheral tissues to the liver and help to lower total serum cholesterol.

**Familial hypercholesterolemia (FH)** is a hereditary disease caused by mutations affecting LDL receptor. Cells from FH individuals have an impaired ability to take up cholesterol via receptor mediated endocytosis. The result of these mutations is a higher than normal level of serum LDL and cholesterol. High LDL levels favor oxidation of their components and ultimately formation of atherosclerotic plaques. Individuals who are homozygous for the disease have very high levels of cholesterol in the blood and usually die of heart disease before age 20. People heterozygous for the disease have higher than normal cholesterol levels and are at high risk for heart attacks and cerebral infarcts.

**Atherosclerosis contributing factors and treatment strategies.** The incidence of atherosclerosis and coronary heart disease correlates with the serum total cholesterol and LDL cholesterol, and is in inverse relationship with HDL concentrations. The major factors affecting the blood lipoprotein levels are heredity, age, gender, nutritional and life style. Regular exercise and the diet rich in fibers, essential fatty acids and other lipotropic factors are effective in lowering plasma LDL and raising HDL. Widely used cholesterol-lowering drugs inhibit HMG-CoA reductase and decrease endogenous cholesterol biosynthesis (statins), or inhibit the absorption of cholesterol and bile acids in intestines (resins, ezetimibe). Other drugs used (fibrates) mainly decrease plasma triacylglycerols and VLDL.



### **Non-protein organic compounds of blood.**

Total nitrogen of the blood includes a protein nitrogen and nonprotein nitrogen (or residual nitrogen):

$$N \text{ total} = N \text{ prot.} + N \text{ res.} \quad N \text{ res.} = 14,3-25 \text{ mmol/L.}$$

**Residual nitrogen** of the blood includes: urea nitrogen (50%), amino acid nitrogen (25%), creatine nitrogen (5%), creatinine nitrogen, ammonia nitrogen, indican nitrogen, bilirubin nitrogen, uric acid nitrogen etc. **Ammonia** level (25-40  $\mu\text{mol/L}$ ) increases in liver diseases, inherited disturbances of ornithine cycle. **Urea** level (3,3-8,3 mmol/L) increases in chronic diseases of kidney, cancer of ureteral ducts, tuberculosis of kidney, some infectious diseases, sepsis and other. Its level decreases in liver diseases (hepatitis, cirrhoses), pregnancy, inherited disturbances of urea cycle. **Creatinine** (53-105  $\mu\text{mol/L}$ ) increases in retentional azotemia, indicates the degree of chronic renal insufficiency. **Uric acid** level (149-405  $\mu\text{mol/L}$ ) increases in gout.

**Azotemia** is the increased level of residual nitrogen in blood.

**Productive azotemia** is observed in an excessive delivery of nitrogenous products to the blood as result of accelerated degradation of tissues proteins in different states: inflammation, wounds, extensive burns, cachexia and other states.

**Retention azotemia** is caused by incomplete urinary discharge of nitrogen containing products on their normal delivery to the blood stream.

1) Renal retention azotemia is caused by reduced excretory function of kidney (reduced renal clearance). Urea is mainly responsible for the increased residual nitrogen level in renal retention azotemia. Urea constitutes 90% of residual nitrogen of blood instead of 50% in normal conditions.

2) Extrarenal retention azotemia may arise from an acute circulatory insufficiency, low arterial pressure, or reduced renal blood flow. Also, the frequent cause of extrarenal retention azotemia is an obstruction to the urine outflow from the kidney.

### **Blood plasma enzymes. Enzyme diagnostics.**

Enzyme diagnostics is one of the branches of enzymology. It has two main directions:

- 1) use of enzymes as reagents for determination of normal and pathological components in serum, urine, gastric juice etc.
- 2) determination of enzyme activity in biological material with a diagnostic purpose.

**Serum enzymes** are divided into 3 groups:

- 1) Cellular enzymes enter the blood from different organs. Their activity in serum depends on enzyme content in organs, molecular weight, intracellular localization, rate of elimination. Cellular enzymes are divided into non-specific and organ specific.
- 2) Secretory enzymes are synthesized by cells, enter the bloodstream and fulfill their specific functions in the circulatory system. These are enzymes of coagulation system and fibrinolysis, choline esterase etc.
- 3) Excretory enzymes are synthesized by glands of GIT and enter the blood (amylase, lipase).

Enzymes synthesis, functioning and breakdown take place continuously and simultaneously; providing their given concentration and activity. Enzymes are localized in different cellular compartments (cytoplasm, lysosomes, cellular membrane, mitochondrions). That is why increased activity of certain enzymes can indicate the degree of severity of cellular damage. Here, we have provided information about enzymes which are most

frequently used in clinical practice for diagnosis, prognosis and therapy monitoring of different pathologies. Their determination in blood serum has high clinical significance.

**$\alpha$ -amylase.** High activity of this enzyme is observed in the liver, skeletal muscles, microvillus of enterocytes, tears, secretion of mammary glands. Pancreas and salivary glands are richest in amylase. Plasma contains two isoenzymes of  $\alpha$ -amylase: pancreatic (P-type) - secreted by pancreas and salivary (S-type) - produced by salivary glands. In norm pancreatic amylase constitutes 40 % of total serum amylase activity, and salivary – 60%. Determination of  $\alpha$ -amylase activity is very important for diagnosis of pancreatic pathology. Two times and more increased activity of  $\alpha$ -amylase strongly indicates pancreatic damage.

In acute pancreatitis,  $\alpha$ -amylase activity in the blood and urine increases 10-30 times. Initial increase of  $\alpha$ -amylase activity is observed within 4-6 hours after the beginning of the disease, reaches peak within 12-24 h; then decreases and returns to norm within 2-6 days. Serum amylase level does not correlate with the degree of severity of pancreatitis. Pathogenetically hyperamylasaemia appears when oedema of interstitial tissue blocks the pancreatic ducts. It characterises fatty pancreatic necrosis. In haemorrhagic pancreatic necrosis, rise in  $\alpha$ -amylase activity is observed in blood with subsequent rapid decrease. This reflects progressing pancreatic necrosis.

**Aminotransferases (ALT and AST).** Aminotransferases catalyze the process of transamination, they are present in every organ and tissue. Isoenzymes of AST are localized both in cytoplasm and in mitochondrions. ALT predominates in cytoplasm. High concentration of AST is noted in heart and skeletal muscles, liver, kidneys, pancreas and erythrocytes. Damage of any of them leads to significant increase of AST in the blood serum. The most significant increase of AST is observed in myocardial damage. In myocardial infarction, AST activity in blood serum can increase 4-5 times. In acute myocardial infarction, 93-98% of patients have high AST activity; the latter has the same dynamic as Creatine Kinase MB (CK-MB). However, CK-MB increase is more significant. Increase in AST activity reveals hepatic pathology. The most significant increase is observed in acute viral and toxic hepatitis. From mild to moderate increase in AST occurs in liver cirrhosis (2-3 times), obstructive jaundice and liver metastasis. It can be also so in skeletal muscular pathology, for example progressive muscular dystrophy; in pancreatitis; intravascular haemolysis.

Low AST activity usually reveals vitamin B6 deficiency, renal failure, pregnancy.

Highest concentration of ALT is noted in the liver cells. Skeletal muscles, kidneys and heart also contain ALT, but much less. Increased ALT activity is most frequently revealed in acute liver and biliary ducts diseases. ALT activity rises significantly in the early stages of acute viral hepatitis: in 50% of patients ALT increases 5 days before jaundice and hepatomegaly appear, in 90% of patients – 2 days before these symptoms. AST/ALT ratio is called **de Ritis ratio**. Its normal value 1-1,3. It decreases in liver diseases and increases in heart diseases. In toxic (alcoholic) liver damage AST activity rises predominantly, where de Ritis ratio exceeds 2. In viral hepatitis de Ritis ratio decreases. This ratio increases in obstructive jaundice, cholecystitis, liver cirrhosis, while ALT and AST activity increase slightly.

**Alkaline phosphatase (ALP).** The isoenzymes of alkaline phosphatase (ALP) are produced by various tissues: intestinal mucous membrane, osteoblasts, biliary ducts, placenta, mammary gland during lactation. This enzyme is situated on the cellular membrane and takes part in transport of phosphorus.

Several isoenzymes of ALP are present in blood serum. Bone, liver and placental ones are the most significant for clinical and diagnostic purposes.

1) Bone ALP. In bones ALP is secreted by osteoblasts. It is possible, that ALP takes part in maturation of a bone matrix and its mineralization. ALP increases with bone formation. Its significant increase in blood serum results from high osteoblastic activity: growth of bones (children show higher ALP activity than adults; it also increases in the last trimester of pregnancy), reactivation after prolonged immobilization, fractures, deforming osteitis, rickets. It is also a characteristic for osteomalacia (malignant bone tumors, multiple myeloma), tuberculosis of bones, leukemia.

2) Liver ALP. There are two isoenzymes. The first one increases in blood serum in biliary obstruction, due to decreased elimination of enzyme with bile. It also increases in pregnancy (the second half). It is the main indicator of biliary tract pathology. The second isoenzyme increases in hepatocellular pathology: viral hepatitis, liver cirrhosis. But this increase is less significant in comparison to aminotransferases. 1/3 patients with jaundice and liver cirrhosis show increase in ALP activity. Rise in ALP activity is also revealed in 20% of patients with primary liver cancer or liver metastasis.

3) Intestinal ALP. It originates from enterocytes, enters the intestinal lumen and is partially absorbed in the blood. It accounts for a small part of total ALP activity. Its activity can be increased in people with I or III blood groups; especially after meals; in intestinal diseases accompanied by diarrhoea.

4) Placental ALP. It normally appears in pregnancy. The highest activity is revealed during the third trimester. It is the most thermostable isoenzyme of ALP. The most significant increase develops in women with eclampsia as a result of placenta damage. Low ALP activity in pregnancy indicates placental insufficiency.

**Gamma-Glutamyl Transpeptidase (GGT).** The determination of GGT is of great significance in diagnosis of hepatic and hepatobiliary pathology. This test is much more sensitive than either ALP or the transaminase test in detecting obstructive jaundice, cholangitis, cholecystitis.

The highest GGT activity is noted in kidneys – 7000 times higher than in blood serum. In healthy individuals serum GGT activity is low. The liver is considered as the main source of normal serum activity, despite the fact that the kidney has the highest level of the enzyme. Pancreas also contains GGT. Small enzyme concentration is detected in intestine, brain, heart, spleen, prostate gland, skeletal muscles. GGT is located in cellular membrane, lysosomes, cytoplasm. Membrane localization of GGT is characteristic for cells with high secretory, excretory or reabsorptional activity. Serum GGT activity increases in any pathology of liver and bile ducts. If GGT activity is normal, liver disease probability is very low. Thus, GGT is good marker for differential diagnosis of liver pathology. The most significant increase is observed in cholestasis and slight increase - in parenchymal liver disease (necrosis of hepatocytes). GGT activity rises on the early stage of the disease and remains high for a long time. Beside this, GGT is a specific indicator of liver disease, because in comparison to ALP its activity is normal in healthy children, pregnant women and patients with bone diseases.

Determination of GGT activity is also used for diagnosis of alcoholic liver disease and its therapy monitoring. Alcohol induces GGT synthesis in the liver and release from cell membranes. It leads to increase of enzyme activity in the blood serum without hepatic cell damage.

GGT test is also used for diagnosis of pancreatic pathology. Nowadays in Europe, this parameter is used even more often than  $\alpha$ -amylase – traditional indicator of pancreatic pathology. 100% of patients with acute pancreatitis show GGT activity 10-20 times higher than normal.

This test can also be useful for laboratory diagnosis of renal pathology. It is proven that GGT activity in urine rises significantly in pyelonephritis, glomerulonephritis and renal calculi. Determination of GGT in urine allows to diagnose the early stages of kidney disease, which is accompanied by proximal renal tubular damage.

**Creatine Kinase (CK).** CK is a dimer and consists of 2 protein subunits: B (brain) and M (muscle), which combine to form 3 isoenzymes:

- 1) CK-BB (CK-1) – brain
- 2) CK-MB (CK-2) – cardiac
- 3) CK-MM (CK-3) – muscle

CK-BB is present in large amount in brain tissue, prostate, stomach, lungs, urinary bladder, urethra, placenta, thyroid gland. CK-MB accounts for 25-46 % of total CK activity in cardiac muscle and less than 5% in skeletal muscle. CK-MM is present mainly in skeletal and cardiac muscle. CK-MM accounts for 94-96 % of total creatine kinase serum activity, CK-MB – 4-6 %, CK-BB – trace amount or is not detectable in serum. Total CK activity increases in different pathologies: traumas, surgical operations, myocardial infarction, reduced perfusion of muscles, myopathy, dermatomyositis, muscular dystrophy, myocarditis, intoxication, hypothyroidism, infectious diseases (typhoid fever). In some cases slight increase occurs in arthritis, congestive heart failure, tachycardia, pulmonary embolism. In myocardial infarction increase in CK activity occurs within 3-6 hours after an onset of pain. However, determination of its activity within 8 hours gives positive results in 31% of cases. CK is a reliable test for myocardial infarction diagnosis within 8-10 hours after onset of pain. It reaches maximal level within 24 hours and returns to norm within the next 48 hours even in extensive myocardial infarction. Relative increase of CK in myocardial infarction is higher than other enzymes. The determination of CK every 4-6 hours during 24 hours is most informative.

CK-MM increases in blood serum in the same cases, as total CK. CK-BB in blood serum slightly or moderately increases in cancer of certain localizations (lungs, intestine, urinary bladder, prostate gland), trauma of cardiac muscle, connecting tissue diseases. During parturition serum CK-BB may be 6 times higher than normal (the source of its activity are uterus and placenta). In neonates and infants the activity is considerably elevated, especially during the first 24 hr post-partum.

**Lactate Dehydrogenase (LDH)** catalyzes reversible reduction of pyruvate to lactate. LDH consists of two subunits – M (muscle) and H (heart). There are 5 isoenzymes in serum, which are distinguished by their subunit composition. They are identified as follows according to decreasing electrophoretic mobility (movement to anode): LDH-1 (H<sub>4</sub>), LDH-2 (H<sub>3</sub>M<sub>1</sub>), LDH-3 (H<sub>2</sub>M<sub>2</sub>), LDH-4 (H<sub>1</sub>M<sub>3</sub>), LDH-5 (M<sub>4</sub>).

LDH is present in cytoplasm of every tissue. In the liver, heart, kidneys, skeletal muscles and erythrocytes LDH activity is 500 times higher than in serum. That is why damage of these organs is accompanied by elevation of serum LDH. Increase in this enzyme occurs in tissue necrosis, especially in acute myocardial injury, haemolysis (erythrocyte damage), injury of kidneys, skeletal muscles, liver, lungs and skin. Significant increase is observed in hemolytic anaemias or B<sub>12</sub>-folate deficiency.

Normal proportion of LDH isoenzymes in serum is: LDH-1 – 15-30%; LDH-2 – 22-50%; LDH-3 – 15-30%; LDH-4 – 0-15%; LDH-5 – 0-15%. In myocardial infarction increase in LDH occurs within 12-32 hours after an onset of pain and remains elevated during 8-14 days. LDH-1 is the most specific test for diagnosis of myocardial infarction. If within 8-24 hr after the onset of pain LDH (and also CK-MB and AST) level does not increase, we can exclude myocardial infarction. Some patients show correlation between LDH level and extensiveness of myocardial injury. In some cases LDH1/LDH2 ratio can give an additional diagnostic information. Its normal range is 0,6-0,7. In acute myocardial infarction it exceeds 1,0 and returns to norm after 2-3 weeks. LDH-1 also increases in tumors of reproductive organs: teratoma, testicle seminoma, ovarian dysgerminoma.

LDH-2, LDH-3 and LDH-4 have an intermediate properties. Activity of these enzymes grows in massive platelet destruction (pulmonary embolism, massive blood transfusions) and lymphatic system involvement. In non-lymphocytic leukemia, LDH-3 and LDH-4 levels increase. LDH-3 also increases in pancreatitis. LDH-4 level rises in viral, toxic and traumatic liver damage, exacerbation of chronic hepatitis, in active phase of rheumatism, in cardiosclerosis, severe diabetes mellitus, acute nephritis, tumors of the liver, prostate, uterine cervix, mammary gland, intestine.

Skeletal muscles, liver, skin, mucous membranes, some kinds of malignant cells contain small amount of LDH-5. Significant increase in LDH-5 occurs in traumas, inflammatory and degenerative muscular diseases and different liver diseases (hepatitis, cirrhosis and others). Oncologic diseases (i.e. lymphocytic leukemia) also can lead to increase in LDH-5. Its activity also grows in active phase of rheumatism, kidney tumors, rejection of kidney transplant, severe diabetes mellitus.

### **Respiratory function of erythrocytes. Transport of O<sub>2</sub> and CO<sub>2</sub>.**

### **Hemoglobin, its synthesis. Metabolism of porphyrins. Metabolism of iron.**

### **Disorders of hemoglobin metabolism: hemoglobinopathy, thalassemia, porphyria.**

Erythrocytes constitute about 44% of the total blood volume (4,5-5x10<sup>12</sup>/L). The life of erythrocytes is 120 days. New synthesized erythrocytes contain ribosomes and elements of endoplasmic reticulum. Mature erythrocytes don't contain ribosomes, mitochondria, lysosomes, Golgi apparatus. Synthesis of erythrocytes is regulated by erythropoietin. Erythropoietin is synthesized in kidney. It is liberated to blood in hypoxia and is transported to bone marrow.

Erythrocytes have unique and relatively simple metabolism:

- 1) Main source of energy is glucose.
- 2) Source of ATP is anaerobic glycolysis.
- 3) The formation of 2,3-bisphosphoglycerate from 1,3-bisphosphoglycerate by Rappoport-Leubering shunt is very important for regulation of affinity of Hb to O<sub>2</sub>.
- 4) 5-10% of glucose are metabolized by means of pentose phosphate pathway. NADPH is necessary for reduction of glutathione. Deficiency of glucose-6-phosphate dehydrogenase is the cause of drug-induced hemolytic anemia
- 5) Glutathione may be synthesized in erythrocytes. It is necessary for elimination of peroxides.
- 6) Autooxidation of Hb results in formation of metHb (1,7% under normal conditions). This is accompanied by formation of O<sub>2</sub><sup>•</sup> (superoxide radical). NADHdependent methemoglobin reductase converts metHb to Hb.


7) The synthesis of glycogen, fatty acids, proteins, nucleic acids does not occur in erythrocytes. Some lipids, for example cholesterol, may be exchanged with corresponding lipids of plasma.

8) Erythrocytes have some enzymes of nucleotide metabolism.

The main function of erythrocytes is the **transport of gases**. Hemoglobin constitutes 95% intracellular proteins of erythrocytes. Hemoglobin is the principal for transport in the blood of both oxygen and CO<sub>2</sub>.

Functions of hemoglobin: respiratory, maintaining acid-base balance (buffer system).

Structure of hemoglobin: prosthetic group (4 hemes), protein part (globin, 4 subunits).



**Heme** is cyclic tetrapyrrole. Tetrapyrroles consist of four molecules of pyrrole linked in a planar ring by four  $\alpha$ -methyl bridges. Tetrapyrrole has 8 substitutions: 4 methyl groups, 2 – vinyl and 2 – propionate ones. One atom of ferrous iron ( $\text{Fe}^{2+}$ ) is at the center of this planar ring.

**Globin** part of HbA<sub>1</sub> is  $\alpha_2\beta_2$  ( $\alpha$  – 141 amino acid residues,  $\beta$  – 146 amino acid residues). Secondary structure: 75% of every polypeptide chain is  $\alpha$ -helix. Tertiary structure is globule. Hydrophobic amino acid radicals are directed inside of protein molecule, hydrophilic amino acid radicals are directed outside. This facilitates the formation of hydrophobic heme pocket. This pocket defends heme iron from oxidation. Quaternary structure: polypeptide chains of Hb are linked by means of hydrophobic interactions and salt bridges. Quaternary structure provides positive cooperative effect between subunits in binding O<sub>2</sub>. Binding last molecule of O<sub>2</sub> occurs 300 times more readily, than the first one. Binding O<sub>2</sub> is accompanied by the rupture of salt bonds between four subunits. Subsequent O<sub>2</sub> binding is facilitated, since it involves a rupture of fewer salt bonds. Iron atoms of deoxyhemoglobin lie about 0,06 nm beyond the plane of the heme ring. On oxygenation the iron atoms move into the plane of the heme ring. This is accompanied by conformational changes and leads to rupture of salt bonds.

Quaternary structure of Hb provides its **allosteric properties**. The affinity of Hb to O<sub>2</sub> is regulated by low molecular mass ligands, for example by CO<sub>2</sub>, 2,3-bisphosphoglycerate (2,3-BPG). 2,3-BPG is formed from the glycolytic intermediate 1,3-bisphosphoglycerate. 2,3-BPG decreases the affinity of Hb to O<sub>2</sub> by forming additional salt bridges.

This plays the important role in adaptive processes in hypoxia, in supplying of embryo by oxygen. 2,3-BPG regulates fetal hemoglobin in less extent, than adult one.

Therefore fetal Hb has a higher affinity to oxygen than does HbA. This provides the transfer of oxygen from Hb of mother to Hb of embryo.

Heterogeneity of Hb		
Ontogenetic	Heterogeneity of adults	Heterogeneity, which is provided by mutations (abnormal Hb)
1. Embryonal HbP is found in first three months of intrauterine life of the baby. Gower I ( $\xi_4$ ) Gower II ( $\alpha_2\xi_2$ ) 2. HbF ( $\alpha_2\gamma_2$ ) 3. HbA ( $\alpha_2\beta_2$ ) Newborns contain 80% HbF and 20% HbA	1. HbA ( $\alpha_2\beta_2$ ) – 90-95%. 2. HbA2 ( $\alpha_2\delta_2$ ) – 2,5%. 3. HbF – 0,5% 4. HbA <sub>1c</sub> (glycosylated): in normal individuals 3-5%, in diabetes mellitus – 6-15%	Examples: HbS (glutamic acid of 6-th position of $\beta$ -chain is replaced by valine). HbM (amino acid sequence is altered either in $\alpha$ - or $\beta$ - chains: or $\alpha_{58}\text{His} \rightarrow \alpha_{58}\text{Tyr}$ , or $\beta_{63}\text{His} \rightarrow \beta_{63}\text{Tyr}$ )

### Derivatives of Hb:

1) **Oxyhemoglobin** (HbO<sub>2</sub>). Oxygen adds to the hemoglobin heme via iron coordination bonds, the iron is in reduced state (Fe<sup>2+</sup>). Factors, influencing the formation of HbO<sub>2</sub>:

- partial pressure of O<sub>2</sub> (P O<sub>2</sub>) favours oxygenation;
- partial pressure of CO<sub>2</sub> (P CO<sub>2</sub>) favours dissociation;
- temperature decreases the affinity of Hb to O<sub>2</sub>;
- pH of the medium (acidosis favours liberation of O<sub>2</sub>);
- 2,3-bisphosphoglycerate diminishes the affinity of Hb to O<sub>2</sub>.

2) **Carboxy-Hb** (HbCO). Oxygen adds to the hemoglobin heme via iron coordination bonds, the iron is in reduced state (Fe<sup>2+</sup>). The affinity of Hb to CO is 210 times more than to O<sub>2</sub>. Dissociation of HbCO is 30 times less than HbO<sub>2</sub>. However, as the partial pressure of oxygen in the inspired air increases, CO is in part eliminated from its binding with hemoglobin.

3) **Carb-Hb** (HbCO<sub>2</sub>). CO<sub>2</sub> combines with NH<sub>2</sub>-group of globin. This is a normal and constant physiologic reaction and accounts for 2 to 10% of CO<sub>2</sub> transported by the blood.


4) **Methemoglobin** (HbOH). It is a derivative in which Fe is in the ferric state (Fe<sup>3+</sup>). In normal healthy adult small amount of methemoglobin may be present (about 1,7% of total Hb). This is converted to normal Hb by methemoglobin reductase.

In vivo HbOH is produced by certain drugs or exposure to certain poisons which are oxidants. Injection of intravenous glucose or methylene blue helps to reduce methemoglobin (Fe<sup>3+</sup>) to Hb (Fe<sup>2+</sup>). Other causes:

- Familial methemoglobinemia is inherited disorder due to lack or absence of the enzyme methemoglobin reductase.
- Methemoglobinemia may also be found in individuals with abnormal hemoglobin as HbM.

The increased formation of HbOH is used for the treatment of cyanide poisoning. The lethal and toxic action of cyanides is provided by inhibition of cytochrome oxidase. Cyanides and HCN do not react directly with hemoglobin but they react with methemoglobin to form cyanmethemoglobin, which is not toxic.

In medical practice of common use is the analysis for blood pigment which is based on a study of spectral properties of hemoglobin heme or of its oxidized products – hemin or hematin, produced by treating hemoglobin with a dilute alkaline solution or with acetic acid in the presence of sodium chloride. Hematin reduced by ammonium sulphite in the presence of globin, produces a hemoglobin derivative hemochromogen. It exhibits a characteristic absorption spectrum. This method is widely used in forensic medical practice for examination of blood spots.



**Figure 8.** Key concept map for hemoglobin structure and function (Harvey, 2011)

### Transport of oxygen.


Oxygen is continuously supplied to the tissue cells. The total requirement of O<sub>2</sub> is around 250 mL/minute in the resting state and more than ten times during vigorous exercise. The requirement of O<sub>2</sub> to the tissue is fulfilled in two ways: oxygen in physical solution; by oxyhemoglobin. A small amount of O<sub>2</sub> can be dissolved to form a solution (0,3 mL of O<sub>2</sub> per 100 mL of blood).

Most of O<sub>2</sub> is supplied to the tissues as HbO<sub>2</sub>. One gram of Hb can carry 1,34 ml of O<sub>2</sub> at complete saturation. The hemoglobin concentration in the blood in healthy individual is 130-160 g/L. At the arterial blood the oxygen saturation of hemoglobin is 96%. Under these conditions the amount of O<sub>2</sub> which is linked with Hb is 19,3 ml of O<sub>2</sub> per 100 ml of blood.



Blood, which contains 150 g/L of Hb:	
Arterial blood	Venous blood
pO <sub>2</sub> 95 mm of Hg	pO <sub>2</sub> 40 mm of Hg
pCO <sub>2</sub> 40 mm of Hg	pCO <sub>2</sub> 46 mm of Hg
Hb: 97% of saturation by O <sub>2</sub>	Hb: 75% of saturation by O <sub>2</sub>


Oxygen supply to tissue cells is facilitated by high pO<sub>2</sub> levels in lungs. It is enhanced by the relatively high pCO<sub>2</sub> (Bohr effect), high acidity (low pH), high temperature in metabolically active tissues.



### Transport of carbon dioxide.

Carbon dioxide is transported from the tissues to lungs at the rate of about 180 ml/min: 6-7% of CO<sub>2</sub> is transported in a physically dissolved state, 3-10% in Hb CO<sub>2</sub>; most CO<sub>2</sub> is transported in the bicarbonate form.


Carbon dioxide is supplied to erythrocytes. It is converted into H<sub>2</sub>CO<sub>3</sub> under the influence of carbonic anhydrase. Deoxyhemoglobin is weak acid, it binds H<sup>+</sup>. Accumulated bicarbonate anions (HCO<sub>3</sub><sup>-</sup>) are transported from erythrocyte cell to plasma. They are exchanged for chloride ions (the chloride shift).



**Figure 9.** Fate of CO<sub>2</sub> in the red blood cell (Janson, 2012)

## Synthesis of hemoproteins.

A specific feature of hemoproteins is the metabolic involvement of the nonprotein moiety of these conjugated proteins. The hemoglobin of blood erythrocytes and of marrow cells accounts for a major portion (about 83%) of hemoproteins in the human organism. The remainder is myoglobin of skeletal muscles and heart (about 17%) and cellular hemoproteins – cytochromes, catalase, and peroxidase (1%).



Glycine and succinyl~S<sub>Co</sub>A are the starting compounds in heme synthesis. The reaction involving the pyridoxal-assisted enzyme δ-aminolevulinic acid synthetase yields δ-aminolevulinic acid. Two molecules of δ-aminolevulinic acid are combined with the participation of porphobilinogen synthetase, to form porphobilinogen, a direct precursor of porphyrins. One of these is coproporphyrin III which is directly converted to protoporphyrin IX. The insertion of iron ions (Fe<sup>2+</sup>) into the protoporphyrin IX ring is effected with the assistance of ferrochelatase. At the ultimate step, heme becomes complexed with globin to form hemoglobin or myoglobin. In the synthesis of other hemoproteins, heme adds to the specific protein moiety of cytochromes or other hemo-containing enzymes. If there is not a sufficient quantity of protein to bind the reaction step of porphyrin synthesis.

**Figure 10.** Heme biosynthesis (Koolman, 2005)

The enzymes involved in heme biosynthesis are found in the marrow, nucleated erythrocytes, liver, kidneys, and intestinal mucosa. The reactions leading to δ-aminolevulinic acid proceed in the mitochondria; the production of porphobilinogen and the subsequent synthesis of coproporphyrinogen III occur in the cytoplasm, and the synthesis of heme from coproporphyrinogen III, in mitochondria.

### **Disturbances of Hb synthesis.**

**Hemoglobinopathies** are due to a hereditary change in the primary structure of peptide chains, for example, in sickle cell anemia HbS ( $\beta$ -chain: 6 Glu  $\rightarrow$  6 Val).

**Thalassemias** are due to a disturbed synthesis of one of the normal hemoglobin chains. For example in  $\beta$ -chain thalassemia an excess of  $\alpha$ -chains occurs which can combine with  $\delta$ -chains producing an increase in HbA<sub>2</sub> or with  $\gamma$ -chains producing an increase in HbF (15-60%).

### **Hereditary Porphyrias:**

1) **Congenital erythropoietic porphyria** (Hunter's disease) is due to a deficiency of uroporphyrinogen III cosynthase. Patients with congenital erythropoietic porphyria excrete large quantities of the type I isomers of both uroporphyrinogen and coproporphyrinogen, which in the urine are spontaneously oxidized to uroporphyrin I and coproporphyrin I, both fluorescent red pigments. Urine is usually red colored. Teeth and bones may be brownish or pink due to porphyrin deposition. Affected individuals exhibit abnormal sensitivity to light (photosensitivity) and development of skin lesions.

2) **Intermittent acute porphyria** (IAP) is due to a deficiency of porphobilinogen deaminase (uroporphyrinogen I synthase). Patients with IAP excrete massive quantities of porphobilinogen and aminolevulinate in the urine. Both of these compounds are colorless, but porphobilinogen upon exposure to light and air polymerizes spontaneously but slowly to form 2 colored compounds: porphobilin and porphyrin. The concentration of  $\delta$ -aminolevulinate and porphobilinogen are increased. They are neurotoxins. Clinical symptoms are abdominal pain, neuropsychiatric symptoms.

3) **Protoporphyrin** is due to a deficiency of ferrochelatase. Photosensitivity is observed.

**Acquired porphyrias** are observed in iron deficiency anemia, in liver diseases, in exposure to toxic compounds.

### **Metabolism of iron.**


Blood uses carrier proteins to transfer essential nutrients as part of each person's metabolism. One important example is iron that is involved in a vast array of important biologic reactions:

1. Binds O<sub>2</sub> as part of Hb molecule. An adult human has approximately 4 g of iron, of which about two-thirds is employed in the O<sub>2</sub>-carrying role of Hb.
2. Mediates a wide variety of oxidative-reductive reactions by serving as an essential cofactor for many proteins via oxidation between ferric (Fe<sup>3+</sup>) and reduced Fe<sup>2+</sup> states.
3. Iron is important for many microorganisms. Keeping iron sequestered from these invaders is an important part of the immune system.


In most well-rounded Western diets, meats and green, leafy vegetables provide adequate iron to prevent iron deficiency. Certain iron-poor diets, such as vegan diets, however, can result in iron deficiency if not supplemented. Once iron is ingested, it is converted from the Fe<sup>3+</sup> state to the Fe<sup>2+</sup> state by intestinal ferric reductase. Fe<sup>2+</sup>, not Fe<sup>3+</sup>, is transported into the intestinal epithelial cell through the divalent metal transporter (DMT1) protein. Fe<sup>2+</sup> is transported out of the intestinal epithelial cells into blood through a second transporter, ferroportin.

Intestinal lumen ferric (Fe<sup>3+</sup>) reductase reduces Fe<sup>3+</sup> to Fe<sup>2+</sup>. Fe<sup>2+</sup> is transported from the lumen into the intestinal epithelial cell through heme transporter (HT), endosomes, and/or divalent metal transporter 1 (DMT1). Fe<sup>2+</sup> can be converted back to Fe<sup>3+</sup> and bound to transferrin within the intestinal cell or can be transported into the blood by ferroportin (FP)

and hephaestin (HP). The  $\text{Fe}^{2+}$  oxidized to  $\text{Fe}^{3+}$ , which binds to plasma transferrin, is carried through the circulation to the tissues.



**Figure 11.** Overview of iron transport (Janson, 2012)



**Figure 12.** Iron transportation by transferrin (Janson, 2012)

This exceedingly high affinity is the biochemical mechanism that the human body has adapted to prevent any free iron from existing in the blood stream. The iron bound transferrin circulates through the blood stream until it binds to a transferrin receptor on the surface of a cell. Cells that express transferrin receptors have high iron demands. These include developing RBCs, dividing cells, and microorganisms. The transferrin-iron-transferrin-receptor complex is endocytosed through the classic clathrin-coated pit pathway. Following endocytosis, pH of the endocytic vesicle is reduced to 5, liberating iron from its carrier protein and making iron available for biologic reactions.

Transferrin transports iron in the blood to the bone marrow to make hemoglobin and red blood cells (erythropoiesis). Transferrin also carries iron to the liver and heart for storage in ferritin molecules, as well as to other parts of the body for various enzymatic and other functions.

**Ferritin.** Although Hb is the most abundant protein that uses iron, ferritin is the most important protein for iron storage. Ferritin consists of a 24-unit multimer of heavy (H) and light (L) chains that create a hollow shell. Iron is imported into the shell as  $\text{Fe}^{2+}$  but is converted to  $\text{Fe}^{3+}$  within the ferritin core by the H chain. A fully loaded molecule of ferritin contains 4500 atoms of iron. It is for this reason that the ferritin molecule is metaphorically referred to as a “bag of rust.” A pathologic form of iron deposition is called **hemosiderin**. This is a nonstructured conglomerate of intracellular iron and is often a pathologic consequence of prolonged inflammation. Prolonged deposition of hemosiderin can cause fibrosis (scarring) of tissues. The importance of iron is highlighted by the fact that the human body has no natural way to excrete it. Unlike other ions, such as sodium, potassium, and calcium, kidneys do not excrete excess iron in the urine. Indeed, the physiological default is to conserve iron. Iron deficiency is much easier to treat than iron overload. Accumulated iron is toxic to a variety of tissues, especially the heart and liver. Infusion of an iron chelator, such as **deferoxamine** or **deferasirox**, provides the only means of reducing some of the iron accumulation. Chelation works by the drug-binding free iron in the blood followed by excretion of the iron-drug complex. The body does lose a small amount of iron each day by the sloughing of skin and epithelial cells and in women through their menstrual flow. However, the same reactivity that makes it a valuable cofactor for proteins also means that free iron can be dangerous because it can easily generate damaging  $\text{O}_2$  free radicals. Thus, the body has a powerful system to control the metabolism of this precious and perilous metal.

**Regulation of iron availability by hepcidin.** Although the human body has no natural way of excreting excess iron, it is able to regulate the uptake and availability of iron through the protein hepcidin. Hepcidin is synthesized as an 84-amino acid precursor, which is then processed to the 25-amino acid active form. Hepcidin acts as a negative regulator of ferroportin, blocking the ability of cells to export iron from the cytoplasm into the blood. In the intestinal cell, this inhibition results in decreased iron absorption from the diet. In reticuloendothelial cells, the primary storage depot for iron, this results in a decreased ability of iron to be mobilized from storage pools to cells that need it. Hepcidin is made in the liver, and the levels of hepcidin in the blood are controlled by a variety of different stimuli, including the total iron stores in the body, the erythropoietic demands of making RBCs, hypoxia, and inflammation. When iron stores are high, the level of hepcidin is increased and the amount of iron absorbed is decreased. When the demand for making RBCs is increased, for example, in response to acute blood loss, the level of hepcidin decreases and the amount of bioavailable iron is increased. Similarly, when tissues do not receive sufficient  $\text{O}_2$  (hypoxia), it signals that more RBCs are needed, and hepcidin production is decreased. Finally, inflammation, such as signaled through the inflammatory cytokine interleukin-6 (IL-6), causes an increase in hepcidin production. This inflammatory regulation is thought to reflect a host defense mechanism because it sequesters iron away from infectious organisms, limiting their growth. If the inflammatory state persists, however, hepcidin sequesters iron away from both the microorganism and the human cells. There is insufficient iron to support the demands for RBC production, and a state of anemia of inflammation or anemia of chronic disease develops.

**Iron deficiency anemia** is the most common nutritional disorder in the world, believed to affect 1 billion people. In children, in the developed world, the most common cause of iron deficiency anemia is the excess consumption of cow’s milk. Excess cow’s milk can cause inflammation damaging the intestinal lining, resulting in blood loss as well as a diminished capacity to absorb iron. The first manifestation of iron deficiency is an anemia that stems from an inadequate iron supply to sustain erythropoiesis. The symptoms of iron

deficiency anemia include pallor, weakness, and lethargy. Severe and prolonged iron deficiency can result in neuropsychological problems. The diagnosis of iron deficiency is usually made by laboratory studies that demonstrate a microcytic anemia (smaller, pale RBCs), reflective of poor Hb production. Blood studies also show low ferritin level and low transferrin saturation (only 10% contain iron as compared with the normal 30-40%). Because ferritin is an acute phase reactant that increases in times of stress and illness, sometimes it can be paradoxically high even in iron-deficient states.

The treatment for iron deficiency is to give iron. The most bioavailable dietary iron is in red meat, but often the patient is unable or unwilling to pursue this method. In this situation, oral elemental iron is given. Because iron is transported by DMT1 as  $Fe^{2+}$ , not  $Fe^{3+}$ , some physicians will advise their patients to simultaneously drink orange juice or take ascorbic acid (vitamin C). Ascorbic acid reduces  $Fe^{3+}$  iron to the  $Fe^{2+}$  state, facilitating the absorption of elemental iron. In treating iron deficiency anemia in children, caused by excessive milk consumption, it is important to both administer elemental iron and dramatically decrease milk consumption. In rare cases of adult and child anemia, such as poor compliance or anatomic or genetic defects in iron absorption, iron is administered intravenously. Supplemental iron is given until the microcytic anemia is resolved and normal levels of ferritin and transferrin saturation are achieved.

**Summary of the causes of some important disorders affecting RBCs (Murray, 2003)**

Disorder	Sole or major cause
Iron deficiency anemia	Inadequate intake or excessive loss of iron
Methemoglobinemia	Intake of excess oxidants (various chemicals and drugs) Genetic deficiency in the NADH-dependent methemoglobin reductase system Inheritance of HbM
Sickle cell anemia	Sequence of codon 6 of the $\beta$ chain changed from GAG in the normal gene to GTG in the sickle cell gene, resulting in substitution of valine for glutamic acid
$\alpha$ -Thalassemias	Mutations in the $\alpha$ -globin genes, mainly unequal crossing-over and large deletions and less commonly nonsense and frameshift mutations
$\beta$ -Thalassemia	A very wide variety of mutations in the $\beta$ -globin gene, including deletions, nonsense and frameshift mutations, and others affecting every aspect of its structure (eg, splice sites, promoter mutants)
Megaloblastic anemias	
Deficiency of vitamin B <sub>12</sub>	Decreased absorption of B <sub>12</sub> , often due to a deficiency of intrinsic factor, normally secreted by gastric parietal cells
Deficiency of folic acid	Decreased intake, defective absorption, or increased demand (eg, in pregnancy) for folate
Hereditary spherocytosis	Deficiencies in the amount or in the structure of $\alpha$ or $\beta$ spectrin, ankyrin, band 3 or band 4.1
Glucose-6-phosphate dehydrogenase (G6PD) deficiency	A variety of mutations in the gene (X-linked) for G6PD, mostly single point mutations
Pyruvate kinase (PK) deficiency	Presumably a variety of mutations in the gene for the R (red cell) isozyme of PK
Paroxysmal nocturnal hemoglobinemia	Mutations in the PIG-A gene, affecting synthesis of GPI-anchored proteins

## CONTROL QUESTIONS

1. Blood as a specialized tissue of the body, its composition. Functions of blood. Blood preparations.
2. Physical and chemical properties of blood. Inorganic components of blood. Imbalance of blood electrolytes (Na, K, Ca).
3. Acid-base balance, its regulation. Buffer blood systems. Acidosis and alkalosis: types, causes, mechanisms of compensation.
4. Blood plasma proteins. Albumins and globulins, their biological role. Hyper-, hypo- and dysproteinemia: causes, clinical symptoms. Paraproteinemia. Acute phase proteins.
5. Blood plasma lipoproteins. Atherosclerosis.
6. Non-protein organic compounds of blood. Residual nitrogen. Azotemia.
7. Blood plasma enzymes. Enzymodiagnosics.
8. Respiratory function of erythrocytes. Hemoglobin (structure, properties). Transport of oxygen and carbon dioxide. Factors affecting the binding of hemoglobin to oxygen. Hemoglobin derivatives.
9. Hemoglobin metabolism, its synthesis. Metabolism of porphyrins. Disorders of hemoglobin metabolism: hemoglobinopathy, thalassemia, porphyria.
10. Metabolism of iron. Iron deficiency anemia.

## LIST OF INFORMATION SOURCES

1. Biological chemistry. Lecture notes / A. D. Tahanovich [et al.]. – Minsk : BSMU, 2017. – 144 p.
2. Biochemistry: Manual for medical students or interns/ L.D. Popova, A.V. Polikarpova – Kharkiv: KNMU, 2011. – 539 p.
3. Gavriliuc Ludmila A. Biochemistry: lectures for students of Medical Departments. – Chisinau, 2011. – 139 p.
4. Janson L.W., Tischler M. Medical Biochemistry: The Big Picture / The McGraw-Hill Companies, 2012. – 431 p.
5. Koolman J. et al. Color Atlas of Biochemistry (2th Ed.) – Thieme, 2005. – 476 p.
6. Lelevich S.V. Clinical biochemistry: manual for students of the Faculty of Foreign Students with English medium of instruction / S.V. Lelevich, T.V. Popechits. – Grodno: GrSMU, 2010. – 86 p.
7. Murray R.K., Granner D.K., Mayes P.A., Rodwell V.W. Harper's illustrated Biochemistry. Twenty-sixth International Edition, 2003. – 702 p.
8. Richard A Harvey, Ph. D.; Denise R Ferrier, Lippincott's Illustrated Reviews: Biochemistry, Fifth Edition. – 2011. – 531 p.


## CHAPTER 2. BIOCHEMISTRY OF LIVER. XENOBIOTICS AND DETOXIFICATION PROCESSES.

### The role of the liver in protein, carbohydrate and lipid metabolism.

#### Biosynthesis of specialized proteins.

#### Digestion, storage and excretion of different metabolites.

Weighing 1,5 kg, the liver is one of the largest organs in the human body. Although it only represents 2-3% of the body mass, it accounts for 25-30% of oxygen consumption. Liver plays the central role in regulation and integration of metabolism. Hepatocytes make up 90% of the cell mass of liver. They are in close contact with the blood, which enters the liver from the portal vein (more than 70%) and the hepatic arteries (30%), flows through capillary vessels known as sinusoids, and is collected again in the central vein of the hepatic lobes. Hepatocytes are particularly rich in endoplasmic reticulum as they carry out intensive protein and lipid synthesis. The cytoplasm contains granules of glycogen. Between hepatocytes there are bile capillaries through which bile components are excreted.



**Figure 13.** Diagram of a hepatocyte (Koolman, 2005)

#### **Functions of liver.**

- 1) Uptake of nutrients supplied by intestine via the portal vein.
- 2) Biosynthesis of endogenous compounds and storage, conversion and degradation of them into excretable molecules (metabolism). In particular, the liver is responsible for the biosynthesis and degradation of almost all plasma proteins.
- 3) Supply of the body with metabolites and nutrients.
- 4) Detoxification of toxic compounds by biotransformation.
- 5) Excretion of substances with the bile.



## Role of the liver in carbohydrate metabolism.

The liver plays the important role in supporting glucose concentration constancy in blood. This is provided by the following mechanisms:

- 1) The liver takes up glucose and other monosaccharides from the plasma.
- 2) Transporters in the plasma membrane of hepatocytes allow insulin-independent transport of glucose and other sugars in both directions. The liver has the enzyme glucokinase, which has higher  $K_m$  (10mM) as compared with hexokinase (0,01-0,1 mM). This enzyme can react by increasing activity in response of the enhance of glucose content in portal vein after food intake.
- 3) Glucose is then either stored in the form of polysaccharide glycogen or converted into fatty acids. When there is a drop of the blood glucose level, the liver releases glucose again by breaking down glycogen. In contrast to muscle, the liver possesses the enzyme glucose-6-phosphatase, which can release glucose from glucose-6-phosphate. Therefore glycogen of liver is used by not only this organ but also by other tissues and organs.
- 4) If the glycogen store is exhausted, glucose also can be synthesized by gluconeogenesis from lactate, glycerol or the carbon skeletons of amino acids. Regeneration of glucose (up to 250 g per day) mainly takes place in the liver. The tubule cells of the kidney are also capable of carrying out gluconeogenesis, but due to their much smaller mass, their contribution only represents around 10% of total glucose formation.
- 5) Fructose and galactose are mainly metabolized by the liver, which channels them into glycolysis.
- 6) The process of glucose utilization is also intensive in liver:
  - metabolites of glycolysis and acetyl-CoA are used for biosynthesis of TAGs;
  - NADPH, which is formed in pentose phosphate pathway, is used for the synthesis of fatty acids and cholesterol;
  - ribose-5-phosphate is used for synthesis of nucleic acids.




Figure 14. Gluconeogenesis in liver (Koolman, 2005)

### Role of liver in lipid metabolism.

The liver is the most important site for the formation of fatty acids, fats (triacylglycerols), cholesterol and the only site for the synthesis of ketone bodies. Most of these products are released into the blood. In contrast, the triacylglycerols synthesized in adipose tissue are stored there.

1) Lipid metabolism in the liver is closely linked with carbohydrates and amino acids metabolism. When it is a good supply of nutrients in the resorptive state, the liver converts glucose via acetyl-CoA into fatty acids. Fatty acids are converted into fats and phospholipids. Together with apoproteins they are packed into VLDLs and then released into the blood by exocytosis.

2) The most of cholesterol (250-500 mg/day, 50-80%) is synthesized in the liver. It is transported to tissues by means of LDLs. LDLs are mainly formed in blood stream from VLDLs. Small amount of LDLs is synthesized immediately in liver. Some cholesterol is required for the synthesis of bile acids. The liver also contributes to the cholesterol metabolism by taking up from the blood and breaking down lipoproteins that contain cholesterol and cholesterol esters (HDLs, IDLs, LDLs).

3) The synthesis of ketone bodies is located in liver. Acetoacetate and  $\beta$ -hydroxybutyrate are alternative fuel for extrahepatic tissues (skeletal muscles, heart, kidneys). Brain also uses ketone bodies in long starvation. Acetone cannot be metabolized and is exhaled via the lungs or excreted with urine.




Figure 15. Lipid metabolism in liver (Koolman, 2005)

### Role of liver in protein metabolism.

- 1) The liver controls the plasma level of amino acids. Excess amino acids are broken down. In patients with severe liver insufficiency level of amino acids in blood is 21mmol/L instead 2,9-4,3 mmol/L.
- 2) The carbon skeleton of amino acids enters intermediary metabolism and serves for glucose synthesis or energy production.
- 3) The liver is the only organ with complete complex of enzymes for urea synthesis. Disturbances of functioning urea cycle lead to accumulation of ammonia in blood and tissues. Brain neurons are the most sensitive to such pathological situation. This is manifested by development of liver encephalopathy and coma.
- 4) Synthesis of choline, creatine, hydroxylation of phenylalanine occurs in the liver.
- 5) Most of the plasma proteins (albumins, 13-18 g/day; 80% of globulins, factors of blood clotting, fibrinolytic systems of blood: II, V, IX, X, XI, XII, XIII fibrinogen, antithrombin, antiplasmin) are synthesized in the liver.

### Storage function.

The liver not only stores energy reserves and nutrients for the body, but also certain mineral substances (including iron) and vitamins (A, D, K, folic acid, B<sub>12</sub>).

**Summary of Liver Functions (Janson. 2012)**

Function	Molecule(s)	Description
Amino acid/ protein metabolism	Amino acid synthesis	Liver enzymes are responsible for most of amino acid synthesis
	Protein degradation	Liver enzymes are responsible for most of protein degradation
Carbohydrate metabolism	Gluconeogenesis	Liver enzymes account for a significant amount of gluconeogenesis
	Glycogenesis	Liver enzymes account for a significant amount of glycogen synthesis
	Glycogenolysis	Liver enzymes account for a significant amount of glycogen breakdown
Lipid metabolism	Lipogenesis	Liver enzymes are responsible for much of lipid (triglyceride) synthesis
	Cholesterol/ lipoproteins	Liver enzymes are responsible for endogenous cholesterol/lipoprotein metabolism. These reactions and pathways are discussed in detail in the text below.
	Apolipoproteins	Major site of <b>apolipoprotein</b> synthesis, proteins responsible for increasing the solubility and transporting dietary fats in the blood, lipoproteins components involved in cholesterol metabolism. Apolipoproteins can also serve as cofactors and can bind to receptors as part of their function.
Coagulation factor synthesis/ clot formation and breakdown	Coagulation factors	Synthesis of coagulation factors I (fibrinogen), II (Prothrombin), V, VII, IX, X, and XI
	Fibronectin (soluble)	Soluble <b>fibronectin</b> , which differs from the insoluble, extracellular matrix form, is a glycoprotein, which, along with <b>fibrin</b> , helps to form the initial blood clot following injury. The soluble fibrin/fibronectin clot is replaced by other matrix proteins, including the insoluble form of fibronectin, as part of the process of wound healing.
	$\alpha_2$ -macroglobulin	Functions as an inhibitor of <b>thrombin</b> coagulation and <b>plasmin/kallikrein</b> fibrinolysis
	$\alpha_1$ -antitrypsin	Serine protease inhibitor, which covalently binds to <b>trypsin</b> and inactivates its function, including cleavage of lung elastase. Deficient $\alpha_1$ -antitrypsin leads to a variety of diseases in the lungs, including <b>cystic fibrosis</b> , and <b>congenital, panacinar emphysema/ chronic obstructive pulmonary disease (COPD)</b> .

	Antithrombin III	Glycoprotein serine protease in blood, which inactivates <b>thrombin</b> (coagulation factor IIa) as well as <b>kallikrein</b> and <b>plasmin</b> molecules, thereby inhibiting clot formation. Antithrombin III activity is increased by the binding of heparin. Antithrombin III also inactivates trypsin and other serine protease enzymes of the classical complement pathway
	Plasminogen/ plasmin	Serine protease, produced as <b>plasminogen</b> in the liver and subsequently converted to <b>plasmin</b> in the blood by <b>coagulation factor XII (Hageman)</b> , <b>tissue plasminogen activator</b> , and/ or <b>urokinase plasminogen activator</b> . Activated plasmin breaks down fibrin/fibronectin clots ( <b>fibrinolysis</b> ). Plasmin activates parts of the complement system and collagen-cleaving enzymes known as <b>collagenases</b> . Self-cleavage of plasminogen produces the molecule <b>angiostatin</b> , a potent inhibitor of the formation of new blood vessels. Plasmin also breaks down the wall of Graafian follicles to allow <b>ovulation</b> .
	$\alpha_2$ -antiplasmin	Serine protease inhibitor, which inactivates <b>plasmin</b> and, thereby, inhibits fibrinolysis, the breakdown of the initial fibrin clot formed upon injury.
<b>Bile synthesis</b>	Bile	Utilized in small intestine for digestion and absorption of lipids. Includes water, bile acids (normally conjugated to taurine or glycine), bile pigments, including bilirubin, from breakdown of hemoglobin porphyrin molecules, cholesterol, phospholipids, and bicarbonate.
<b>Breakdown of hemoglobin</b>	Bilirubin	Breakdown of heme from degenerating red blood cells starts in the spleen with reduction of the heme by reduced nicotinamide adenine dinucleotide phosphate (NADPH) to biliverdin/free $Fe^{3+}$ ion and then a further reduction with NADPH to bilirubin. This form is known as <b>unconjugated bilirubin</b> . Bilirubin is transported by albumin (see below) to the liver where it is conjugated to glucuronic acid by the enzyme <b>UDP-glucuronosyltransferase</b> , a process that makes the molecule more soluble. This form is known as <b>conjugated bilirubin</b> and is excreted in bile into the intestine where it is converted by bacteria into <b>urobilinogen</b> . Urobilinogens are partly absorbed in the intestine and finally excreted in urine as <b>urobilin</b> or are converted to <b>stercobilin</b> for excretion in feces. Urobilinogens are responsible for the yellow color of urine; stercobilin is responsible for the brown color of feces.
<b>Urea cycle</b>	Urea	Main site of conversion of amino acid nitrogen to urea via the urea cycle. Urea synthesis also occurs, although to a lesser extent, in the kidney.
<b>Detoxification</b>	Various toxins, drugs, and alcohol	Breakdown and elimination of a variety of toxic substances, including <b>toxins</b> , <b>medications</b> , and <b>alcohol</b> . Many toxic molecules are normally metabolized in two overall steps (phases I and II). Drug detoxification also occurs to a lesser extent in the digestive system, lungs, kidneys, and skin. <b>Phase I</b> normally occurs prior to phase II reactions and usually involves reactions that increase the polar nature of the molecule (e.g., reduction/oxidation, hydrolysis, and cyclization/decyclization). Many drugs are designed to be activated, inactivated, or modified for elimination in urine or feces (via bile conjugation) by phase I reactions. Phase I enzymes in the liver include the <b>cytochrome p450 system</b> (oxidation and reduction reactions) and alcohol dehydrogenase (converts alcohol to acetaldehyde) and acetaldehyde dehydrogenase (converts acetaldehyde to acetic acid). Alcohol metabolism can also occur in other tissues, including stomach epithelium (men only) and the brain. <b>Phase II</b> reactions normally involve the addition of biochemical groups (e.g., glucuronic acid, sulfonates, glutathione, methylation, acetylation, and/or amino acid residues) to polar groups added in phase I, including carboxyl (COOH), hydroxyl (OH), amino (NH <sub>2</sub> ), and sulfhydryl (SH) groups. Phase II reactions normally permanently inactivate the toxin or drug.
<b>Storage</b>	Albumin	Important carrier protein of multiple molecules in the blood, including thyroid hormones and other fat-soluble hormones (see additional hormone transport proteins below). Also, transports fatty acids on their way to the liver for storage, or oxidation for energy generation, unconjugated bilirubin (see above), and several medications. Other globulins produced by the liver serve the same role(s), although to a lesser extent than albumin.
	Glycogen	The liver stores glucose in the form of <b>glycogen</b>

	Vitamin A (retinol)	Stores vitamin A transported from the intestine esterified with palmitate via chylomicrons. The liver can store up to a 2-year supply of vitamin A. See below for release, transport, and use.
	Vitamin B <sub>12</sub> (cobalamin)	Storage of vitamin B <sub>12</sub> (approximately 50% of body's total). Because of the efficient recirculation and restorage by the liver, years worth of vitamin B <sub>12</sub> can be stored.
	Vitamin D (as calcidiol)	Vitamin D in the liver is converted by carbon 25-hydroxylation of vitamin D <sub>3</sub> ( <b>cholecalciferol</b> ) by cholecalciferol 25-hydroxylase into a prohormone form called " <b>calcidiol</b> " (25-hydroxy vitamin D <sub>3</sub> ). When released for use, calcidiol is then converted to its active form by a second hydroxylation at the 1 position to form " <b>calcitriol</b> " (1, 25-dihydroxy vitamin D <sub>3</sub> ) by the kidney. The liver can store up to a 4-month supply.
	Vitamin E	Ingested vitamin E is taken up by the liver but only the $\alpha$ -tocopherol form is stored. Other forms ( $\beta$ -, $\gamma$ -, and $\delta$ -tocopherols and $\alpha$ -, $\beta$ -, $\gamma$ -, and $\delta$ -tocotrienols) are metabolized and excreted.
<b>Transport/ "carrier" proteins</b>	$\alpha$ -fetoprotein	Binds to calcium ions affecting the total, available calcium concentration. Serves as a pH buffer and as an osmotic molecule to maintain colloid osmotic pressure (oncotic pressure) that influences the movement of water from and to the blood. Serves a similar role in the developing fetus and, as such, is used as part of a prenatal screen for Down syndrome, neural tube defects, and abdominal wall defects (omphalocele). Serves as a tumor marker for cancer of liver cells and some germ cell and testicular cancers.
	Ceruloplasmin	Enzyme that contains six copper atoms and is responsible for carrying approximately 90% of the body's total copper (additional 10% is contained in albumin).
	Haptoglobin	Transports free hemoglobin molecules released from degenerating red blood cells. Also produced by several other tissues, including kidney, skin, and lung.
	Hemopexin	Transports free heme porphyrin molecule released from degrading hemoglobin. In doing so, it preserves iron and protects the body from the damaging oxidative effects of the free heme group.
	Insulin-like growth factor 1 (IGF-1)-binding protein	Transports IGF-1
	Retinol-binding protein	Binds to de-esterified, alcohol form of retinol (vitamin A) released from storage in the liver and transports to tissues in the body.
	Sex hormonebinding protein	Transports testosterone and estradiol. Also produced in the placenta, testes, uterus, and brain.
	Thyroxin-binding globulin	Transports thyroxine (T <sub>4</sub> ) and 3,5,3'-triiodothyronine (T <sub>3</sub> ).
	Transcortin	Transports cortisol, aldosterone, and progesterone.
	Transferrin	Important carrier protein of iron (Fe <sup>3+</sup> ) as well as the primary store of iron in the body. Transferrin is composed of two, identical monomers, linked by disulfide bonds, each of which can bind and carry one or two Fe <sup>3+</sup> ions. Primarily produced in liver but also made in other tissues (e.g., brain).
	Transthyretin	Transports thyroxine (T <sub>4</sub> ). Also produced in the choroid plexus and retinal pigment epithelium.
Vitamin D-binding protein	Transports vitamin D to tissues in the body. Also, has actin binding activity that may serve as a scavenger role for actin monomers released from injured cells or tissues.	
<b>Miscellaneous</b>	Angiotensinogen	Peptide hormone that, when converted to angiotensin I by the enzyme renin (and subsequently to angiotensin II by angiotensin-converting enzyme), raises blood pressure via a number of mechanisms

C-reactive protein (CRP)	Protein whose liver production is increased because of inflammation, specifically the release of interleukin 6 (IL-6) by macrophages and adipocytes. CRP binds to phosphocholine molecules on degenerating cells to activate the complement system, leading to their phagocytosis by macrophages; CRP may play other roles in the immune system. CRP is used as a marker for inflammation. Its use for risk assessment for heart attack, high blood pressure, high cholesterol/lipids, and diabetes is still being investigated but has been shown not to be as useful as once thought. Investigations of CRP measurement for cancer screening are also ongoing.
Complement proteins	Synthesis of complement proteins C 1–9, including the complement component 3 utilized in both the classical and alternative complement pathways
Insulin	The majority of insulin is degraded in liver cells. Other cells are also able to breakdown insulin as well as other hormones.
IGF-1	Polypeptide hormone mainly responsible for growth in early childhood. Also produced as an autocrine hormone in several target tissues
Fetal red blood cell (erythrocyte) production	Site of production of fetal red blood cells (erythrocytes), containing hemoglobin F. The liver is the sole site of production during the first trimester and is gradually replaced by the developing bone marrow.
Kupffer cells (reticuloendothelial system)	Monocyte/macrophage-type cells of the reticuloendothelial system, which serve as antigen monitors sampling circulating antigens to determine whether an immune response should be mounted.
Thrombopoietin	Glycoprotein hormone that promotes the production of platelets by bone marrow from precursor megakaryocyte cells. Also produced in the kidney.

### **Hemoglobin metabolism, its breakdown. Bile formation.**

The life of red cells is 120 days. After this period of time, the erythrocytes suffer degradation to release hemoglobin. About 6g Hb are renovated every day in adult human. This leads to releasing about 25 mg of iron. Free iron is toxic, but association with transferrin diminishes its potential toxicity and also directs iron to where it is required in the body. The major organs responsible for the erythrocytolysis and hemoglobin breakdown are: liver, spleen, bone marrow.

The **catabolism of heme** from all the heme proteins is carried out in the microsomal fraction of reticuloendothelial cells by a complex enzyme system called heme oxygenase. This process starts with a cleavage of the  $\alpha$ -methylene bridge between the rings I and II of the porphyrin ring system to produce verdoglobin and carbon monoxide. Verdoglobin is spontaneously cleft into iron (ferric ion), globin and biliverdin. Biliverdin reductase reduces the methenyl bridge between pyrrole III and pyrrole IV to a methylene group to produce bilirubin. The chemical conversion of heme to bilirubin by the reticuloendothelial cells can be observed in vivo as the purple color of the heme in a hematoma is slowly converted to the yellow pigment of bilirubin. The further metabolism of bilirubin occurs primarily in the liver. It can be divided into 3 processes:


- 1) Uptake of bilirubin by liver parenchymal cells.
- 2) Conjugation of bilirubin in the smooth endoplasmic reticulum.
- 3) Secretion of conjugated bilirubin into the bile.

**Bilirubin** is hydrophobic molecule. Therefore bilirubin formed in cells of reticulo-endothelial system of spleen and bone marrow is transported to the liver by plasma albumin. Each molecule of albumin appears to have one high affinity site and one low affinity site for bilirubin. In 100 ml of plasma approximately 25 mg of bilirubin can be tightly bound to albumin. 1g of Hb degradation yields 35 mg of bilirubin. The daily bilirubin formation in human adults is approximately 250-350 mg, deriving mainly from Hb, but also from various other hemoproteins such as cytochrome P<sub>450</sub>. Bilirubin which is not bound with albumin, is

toxic, because it is hydrophobic. It can pass through blood-brain barrier and can form complexes with collagen of intercellular matrix and with lipids of membranes. The diminishing pH of blood decreases the affinity of albumin to bilirubin. Some drugs compete with bilirubin to bind with high affinity site of albumin.

In the liver, the bilirubin is removed from albumin and taken up by hepatocytes with a carrier-mediated saturable system. In smooth endoplasmic reticulum of hepatocytes bilirubin is converted to a polar form by adding glucuronic acid molecules to it. This process is called conjugation and is catalyzed by **UDP-glucuronyl transferase**. Secretion of conjugated bilirubin into the bile occurs by an active transport mechanism. Secretion of bilirubin is the rate limiting stage of metabolism of bilirubin in liver. Conjugation and secretion of bilirubin are induced by phenobarbital.


Cells of reticulo-endothelial system of liver, spleen, bone marrow	Hemoglobin	
	↓	+ 2NADPH <sub>2</sub> +O <sub>2</sub>
	Verdoglobin	<b>heme oxygenase</b>
	↓	- Fe <sup>3+</sup> +globin
	biliverdin (green pigment)	<b>bilirubin reductase</b>
	↓	
	bilirubin	
	↓	
Blood	bilirubin + albumin	indirect, free, unconjugated
	↓	
	bilirubin	
	↓	+ UDP-glucuronic acid
Liver	bilirubin monoglucuronide (30%) + bilirubin diglucuronide (70%)	<b>UDP-glucuronyl transferase</b> direct, conjugated
	↓	
Bile	bilirubin glucuronides	
	↓	
	bilirubin glucuronide	
	↓	<b>β-glucuronidase</b> (bacterial enzyme)
Intestine: terminal ileum and the large intestine	bilirubin	
	↓	
	stercobilinogen (urobilinogen)	
	↓	
	stercobilin (feces) 40-280 mg/day urobilin (urine) 4 mg/day	



**Figure 16.** Structure of bilirubin diglucuronide (Murray, 2003)

Small amount of conjugated bilirubin enters to blood. This bilirubin is called direct, conjugated. Direct bilirubin is hydrophilic, therefore only direct bilirubin may be excreted in the urine. As the conjugated bilirubin reaches the terminal ileum and the large intestine, the glucuronides are removed by specific bacterial enzymes ( $\beta$ -glucuronidases). Then bilirubin is finally reduced to stercobilinogen (urobilinogen) by influence of microflora enzymes. Part of it is absorbed, supplies to liver and then or is excreted with bile (large amount) or enters to blood and is excreted in the urine (small amount). Stercobilinogen, which is excreted with feces, is oxidized to stercobilin.

**Total bilirubin** of blood – 8,5-20  $\mu\text{mol/L}$  (75% indirect 25% direct). Determination of bile pigments is very important for differential diagnosis of different forms of jaundice.




**Figure 17. Bilirubin metabolism (Koolman, 2005)**

### **Bile formation function of liver.**

Bile is an important product released by hepatocytes. It promotes the digestion of fats from food by emulsifying them in small intestine. The emulsifying components of bile, apart from phospholipids, mainly consist of bile acids and bile salts. The bile also contains bile pigments, free cholesterol, which are excreted by this way. The cholesterol excreted with the bile is poorly water-soluble. Together with phospholipids and bile acids it forms micelles, which keep it in solution. If the proportions of phospholipids, bile acids and cholesterol shift, gallstones can arise. These mainly consist of precipitated cholesterol (cholesterol stones), but can also contain  $\text{Ca}^{2+}$  salts of bile acids and bile pigments.





**Figure 18.** Metabolism of bile salts (Koolman, 2005)

**Biochemistry of jaundice (hemolytic, hepatic, obstructive): causes, clinical symptoms, differential diagnostics.**

**Hyperbilirubinemia causes jaundice**

When bilirubin in the blood exceeds 1 mg/dL (17.1  $\mu\text{mol/L}$ ), hyperbilirubinemia exists. Hyperbilirubinemia may be due to the production of more bilirubin than the normal liver can excrete, or it may result from the failure of a damaged liver to excrete bilirubin produced in normal amounts. In the absence of hepatic damage, obstruction of the excretory ducts of the liver – by preventing the excretion of bilirubin – will also cause hyperbilirubinemia. In all these situations, bilirubin accumulates in the blood, and when it reaches a certain concentration (approximately 2-2.5 mg/dL), it diffuses into the tissues, which then become yellow. That condition is called **jaundice** or **icterus**.

In clinical studies of jaundice, measurement of bilirubin in the serum is of great value. A method for quantitatively assaying the bilirubin content of the serum was first devised by van den Bergh by application of Ehrlich's test for bilirubin in urine. The Ehrlich reaction is based on the coupling of diazotized sulfanilic acid (Ehrlich's diazo reagent) and bilirubin to produce a reddish-purple azo compound. In the original procedure as described by Ehrlich, methanol was used to provide a solution in which both bilirubin and the diazo reagent were soluble. Van den Bergh inadvertently omitted the methanol on an occasion when assay of bile pigment in human bile was being attempted. To his surprise, normal development of the color occurred "directly." This form of bilirubin that would react without the addition of methanol was thus termed "**directreacting.**"

It was then found that this same direct reaction would also occur in serum from cases of jaundice due to biliary obstruction. However, it was still necessary to add methanol to

detect bilirubin in normal serum or that which was present in excess in serum from cases of hemolytic jaundice where no evidence of obstruction was to be found. To that form of bilirubin which could be measured only after the addition of methanol, the term “**indirect-reacting**” was applied. It was subsequently discovered that the indirect bilirubin is “free” (unconjugated) bilirubin en route to the liver from the reticuloendothelial tissues, where the bilirubin was originally produced by the breakdown of heme porphyrins. Since this bilirubin is not water-soluble, it requires methanol to initiate coupling with the diazo reagent. In the liver, the free bilirubin becomes conjugated with glucuronic acid, and the conjugate, bilirubin glucuronide, can then be excreted into the bile. Furthermore, conjugated bilirubin, being water-soluble, can react directly with the diazo reagent, so that the “direct bilirubin” of van den Bergh is actually a bilirubin conjugate (bilirubin glucuronide).

Depending on the type of bilirubin present in plasma – ie, unconjugated or conjugated – hyperbilirubinemia may be classified as **retention hyperbilirubinemia**, due to overproduction, or **regurgitation hyperbilirubinemia**, due to reflux into the bloodstream because of biliary obstruction.

Because of its hydrophobicity, only unconjugated bilirubin can cross the blood-brain barrier into the central nervous system; thus, encephalopathy due to hyperbilirubinemia (**kernicterus**) can occur only in connection with unconjugated bilirubin, as found in retention hyperbilirubinemia. On the other hand, because of its water-solubility, only conjugated bilirubin can appear in urine. Accordingly, **choluric jaundice** (choluria is the presence of bile pigments in the urine) occurs only in regurgitation hyperbilirubinemia, and **acholuric jaundice** occurs only in the presence of an excess of unconjugated bilirubin.

#### **Elevated amounts of unconjugated bilirubin in blood occur in a number of conditions**

**Hemolytic anemias.** Hemolytic anemias are important causes of unconjugated hyperbilirubinemia, though unconjugated hyperbilirubinemia is usually only slight (<4 mg/dL; <68.4  $\mu\text{mol/L}$ ) even in the event of extensive hemolysis because of the healthy liver’s large capacity for handling bilirubin.

**Neonatal “physiologic jaundice”.** This transient condition is the most common cause of unconjugated hyperbilirubinemia. It results from an accelerated hemolysis around the time of birth and an immature hepatic system for the uptake, conjugation, and secretion of bilirubin. Not only is the bilirubin-UGT activity reduced, but there probably is reduced synthesis of the substrate for that enzyme, UDP-glucuronic acid. Since the increased amount of bilirubin is unconjugated, it is capable of penetrating the blood-brain barrier when its concentration in plasma exceeds that which can be tightly bound by albumin (20-25 mg/dL). This can result in a hyperbilirubinemic toxic encephalopathy, or **kernicterus**, which can cause mental retardation. Because of the recognized inducibility of this bilirubin-metabolizing system, phenobarbital has been administered to jaundiced neonates and is effective in this disorder. In addition, exposure to blue light (phototherapy) promotes the hepatic excretion of unconjugated bilirubin by converting some of the bilirubin to other derivatives such as maleimide fragments and geometric isomers that are excreted in the bile.

**Crigler-Najjar syndrome, type I; congenital nonhemolytic jaundice.** Type I Crigler-Najjar syndrome is a rare autosomal recessive disorder. It is characterized by severe congenital jaundice (serum bilirubin usually exceeds 20 mg/dL) due to mutations in the gene encoding bilirubin-UGT activity in hepatic tissues. The disease is often fatal within the first 15 months of life. Children with this condition have been treated with phototherapy, resulting in some reduction in plasma bilirubin levels. Phenobarbital has no effect on the

formation of bilirubin glucuronides in patients with type I Crigler-Najjar syndrome. A liver transplant may be curative.

**Crigler-Najjar syndrome, type II.** This rare inherited disorder also results from mutations in the gene encoding bilirubin-UGT, but some activity of the enzyme is retained and the condition has a more benign course than type I. Serum bilirubin concentrations usually do not exceed 20 mg/dL. Patients with this condition can respond to treatment with large doses of phenobarbital.

**Gilbert syndrome.** Again, this is caused by mutations in the gene encoding bilirubin-UGT, but approximately 30% of the enzyme's activity is preserved and the condition is entirely harmless.

**Toxic hyperbilirubinemia.** Unconjugated hyperbilirubinemia can result from toxin-induced liver dysfunction such as that caused by chloroform, arsphenamines, carbon tetrachloride, acetaminophen, hepatitis virus, cirrhosis, and Amanita mushroom poisoning. These acquired disorders are due to hepatic parenchymal cell damage, which impairs conjugation.

### **Obstruction in biliary tree is the commonest cause of conjugated hyperbilirubinemia**

**Obstruction of the biliary tree.** Conjugated hyperbilirubinemia commonly results from blockage of the hepatic or common bile ducts, most often due to a gallstone or to cancer of the head of the pancreas. Because of the obstruction, bilirubin diglucuronide cannot be excreted. It thus regurgitates into the hepatic veins and lymphatics, and conjugated bilirubin appears in the blood and urine (choluric jaundice).

The term **cholestatic jaundice** is used to include all cases of extrahepatic obstructive jaundice. It also covers those cases of jaundice that exhibit conjugated hyperbilirubinemia due to micro-obstruction of intrahepatic biliary ductules by swollen, damaged hepatocytes (eg, as may occur in infectious hepatitis).

**Dubin-Johnson syndrome.** This benign autosomal recessive disorder consists of conjugated hyperbilirubinemia in childhood or during adult life. The hyperbilirubinemia is caused by mutations in the gene encoding MRP-2, the protein involved in the secretion of conjugated bilirubin into bile. The centrilobular hepatocytes contain an abnormal black pigment that may be derived from epinephrine.

**Rotor syndrome.** This is a rare benign condition characterized by chronic conjugated hyperbilirubinemia and normal liver histology. Its precise cause has not been identified, but it is thought to be due to an abnormality in hepatic storage.

### **Some conjugated bilirubin can bind covalently to albumin.**

When levels of conjugated bilirubin remain high in plasma, a fraction can bind covalently to albumin (delta bilirubin). Because it is bound covalently to albumin, this fraction has a longer half-life in plasma than does conventional conjugated bilirubin. Thus, it remains elevated during the recovery phase of obstructive jaundice after the remainder of the conjugated bilirubin has declined to normal levels; this explains why some patients continue to appear jaundiced after conjugated bilirubin levels have returned to normal.

### **Urobilinogen and bilirubin in urine are clinical indicators.**

Normally, there are mere traces of urobilinogen in the urine. In **complete obstruction of the bile duct**, no urobilinogen is found in the urine, since bilirubin has no access to the intestine, where it can be converted to urobilinogen. In this case, the presence of bilirubin (conjugated) in the urine without urobilinogen suggests obstructive jaundice, either intrahepatic or posthepatic.

In **jaundice secondary to hemolysis**, the increased production of bilirubin leads to increased production of urobilinogen, which appears in the urine in large amounts. Bilirubin is not usually found in the urine in hemolytic jaundice (because unconjugated bilirubin does not pass into the urine), so that the combination of increased urobilinogen and absence of bilirubin is suggestive of hemolytic jaundice. Increased blood destruction from any cause brings about an increase in urine urobilinogen.

Condition	Serum Bilirubin	Urine Urobilinogen	Urine Bilirubin	Fecal Urobilinogen
Normal	Direct: 0.1–0.4 mg/dL Indirect: 0.2–0.7 mg/dL	0–4 mg/24 h	Absent	40–280 mg/24 h
Hemolytic anemia Hepatitis	↑ Indirect ↑ Direct and indirect	Increased Decreased if micro-obstruction is present	Absent Present if micro-obstruction occurs	Increased Decreased
Obstructive jaundice <sup>1</sup>	↑ Direct	Absent	Present	Trace to absent

**Figure 19.** Laboratory results in normal and patients with three different causes of jaundice (Murray, 2003)

The commonest causes of obstructive (posthepatic) jaundice are cancer of the head of the pancreas and a gallstone lodged in the common bile duct. The presence of bilirubin in the urine is sometimes referred to as choluria – therefore, hepatitis and obstruction of the common bile duct cause choluric jaundice, whereas the jaundice of hemolytic anemia is referred to as acholuric. The laboratory results in patients with hepatitis are variable, depending on the extent of damage to parenchymal cells and the extent of micro-obstruction to bile ductules. Serum levels of ALT and AST are usually markedly elevated in hepatitis, whereas serum levels of alkaline phosphatase are elevated in obstructive liver disease.


Figure 7 summarizes laboratory results obtained on patients with three different causes of jaundice – hemolytic anemia (a prehepatic cause), hepatitis (a hepatic cause), and obstruction of the common bile duct (a posthepatic cause). Laboratory tests on blood (evaluation of the possibility of a hemolytic anemia and measurement of prothrombin time) and on serum (eg, electrophoresis of proteins; activities of the enzymes ALT, AST, and alkaline phosphatase) are also important in helping to distinguish between prehepatic, hepatic, and posthepatic causes of jaundice.

### **Biotransformation of xenobiotics and endogenous toxins. Microsomal oxidation.**

All the substances supplied to the organism in a variety of ways pass through several basically similar stages such as absorption, distribution (mechanical transport), and excretion. The transit rate of substance at these stages may either be increased, or lowered, depending on the structural features and physico-chemical properties of a substance as well as on its affinity to biological molecules. The discipline, dealing with rate characteristics at the stages in which any substance entering the organism is involved, is referred to as chemobiokinetics which treats, in a broader sense, movements of substances in the living organism. Conceptually, chemobiokinetics is divided into three subdisciplines: pharmacokinetics, toxicokinetics, and biokinetics. Pharmacokinetics confines itself to the study of drugs; toxicokinetics, to the study of toxic substances; and biokinetics, to the study of substances not alien to the organism. In many respects, this classification is rather arbitrary, since the distinction between a drug and a poison in many instances may be

evasive. Moreover, even autogenous compounds taken in improper doses may exhibit toxic properties. The subsequent history of a substance after its uptake by the organism is dependent to a significant degree on the rates at which it is converted by various enzymes, i.e. on its metabolic transformations. In point of fact, the metabolism of biogenous substances and xenobiotics used as drugs is governed by the laws of enzymic kinetics. Biogenous substances, being natural substrates for enzymes, are converted at the rates characteristic of catalytic properties of the enzymes involved. The metabolic evolution of xenobiotics is dependent on the occurrence of enzymes capable of catalyzing the conversion of these xenobiotics. If no enzymes that are potentially capable of catalytic intervention of the xenobiotics are available, the xenobiotics behave as metabolically inert. Apparently, in the course of evolution, highly substrate-specific enzymes have laid a basis for the intrinsic metabolism in living organisms, while the enzymes with low specificity towards substrates have taken up defense functions aimed at the inactivation of extraneous invaders.

Biochemistry studies enzyme-assisted conversions of drugs in the organism by making use of appropriate methods and techniques. The drug metabolism in the organism may be represented within the framework of a general scheme:



The drug metabolism is studied by determining the drugs and their metabolites in biological fluids, tissues, and excretions as well as by estimating the activity and kinetics of enzymes involved in the drug metabolism.

Experimentally, the two approaches are used in the studies on metabolism of xenobiotics. In the clinic, the drug metabolism is assessed, as a rule, by measuring the concentration of administered drug and its metabolites in blood, urine, and other excretions.


### **Stages in the metabolism of xenobiotics.**

Biogenous substances, as distinct from xenobiotics, are involved in the conventional metabolic process. Xenobiotics, in the course of their conversion, are subject to two major stages: modification (nonsynthetic stage) and conjugation (synthetic stage).

The **modification stage** is an enzyme-assisted modification of the initial structure of a xenobiotic resulting either in a cleavage of bonds within the xenobiotic molecule, or in the insertion of additional functional groups (e.g. hydroxyl or amino groups) into its molecule, or in a release of its functional groups blocked in the initial structure (for example, by hydrolysis of ester or peptide bonds). The modification leads to an increased solubility of the xenobiotic (xenobiotic becomes more hydrophilic). Additional functional groups are needed to enable the xenobiotic to enter the conjugation stage.

**Conjugation stage** is viewed as an enzyme-assisted process for building covalent bonds between the xenobiotic and biomolecules occurring in the organism's media (e.g. glucuronic acid, sulphates, and others). The conjugation stage terminates in the synthesis of a novel compound whose constituents are, on the one hand, the xenobiotic moiety and, on the other hand, a conjugate (biomolecule).

**Relationship between metabolism of xenobiotics and their structure.** Xenobiotics invaded into the organism are liable to a chain of modifications, or nonsynthetic conversions (oxidation-reduction, isomerization, cyclization, ring opening, and hydrolysis) carried out by the respective enzymes (oxidoreductases, isomerases, lyases, and hydrolases):



Depending on the number of functional groups in the molecule of modified xenobiotic, its conjugation can proceed by a variety of routes in which each of the xenobiotic functional groups becomes bound with a conjugating agent. If the xenobiotic is not functionalized (e.g. benzene), it cannot enter the conjugation stage. In contrast, if the introduced xenobiotic is in possession of an appropriate functional group (e.g., 4-aminobiphenyl); it may become immediately engaged in the conjugation stage with UDP-glucuronic acid.

The knowledge of principles that govern the enzyme-assisted conversions of xenobiotics provides an opportunity to prognosticate metabolic behaviour of any xenobiotic taking into account its structural specificities.

### **Xenobiotic routes in the organism.**

Xenobiotics are either eliminated from the organism, or become accumulated in tissues. Xenobiotics are excreted as:

- 1) supplied (unmodified by enzymes);
- 2) metabolites (modified by enzymes);
- 3) conjugates (by action of conjugating enzymes);
- 4) complexed with biomolecules (for example, metal-containing xenobiotics become bound to cysteine by glutathione and excreted as complexes).

The xenobiotics that accumulate in the organism are those capable of interacting with macromolecules (proteins, nucleic acids, and lipid entities). For example, organochloric compounds, which are readily soluble in lipids, are quite resistant to catabolic conversion and are difficult to eliminate from the organism. They tend to accumulate in lipid-rich tissues. Heavy metals (mercury, cadmium, silver, arsenic, and lead) and preparations containing organometallic compounds become bound with proteins and likewise accumulate in the organism.

### **Metabolism and physiological action of drugs.**

Substances introduced into organism may exhibit either medicinal or toxic properties. Commonly, any drug can exert both medicating and side (toxic) effects. Therefore, generally speaking, the more active the drug, the faster its toxic properties become manifest. During metabolism, the specific activity and toxicity of xenobiotics are susceptible to alterations. Biological activity alterations show up in:

- 1) deactivation, i.e. a loss of medicinal or biological activity of drugs;
- 2) activation, i.e. induced activity of an inactive preparation;
- 3) modification of the major effect, i.e. when the administered drug, on having metabolized, exhibits properties different from those of the initial preparation.

The alterations of toxicity are manifested in:

- 1) deintoxication, i.e. a loss or reduced toxicity of drug;
- 2) toxification, i.e. enhanced toxicity of drug.

The above instances may be exemplified as follows:

**Deactivation** is observed as the functional groups responsible for the biological activity of a drug are either eliminated from, or blocked in the drug molecule. For example, the active sulphanilamide, after its conjugation with acetyl-ScoA, is converted to an inactive acetylsulphanilamide.

**Activation** is observed when the biologically active groups that have been blocked in the initial preparation become deblocked during metabolism:

**Phthalsulphathiazole (inactive drug) → hydrolysis → Sulphathiazole (active drug) + Phthalic acid**

or acquire functional groups that are necessary for eliciting the drug activity:

**Benzopyrene (inactive procarcinogen) → hydrolysis → Hydroxybenzopyrene (carcinogen)**

**Modification** of the major drug effect manifests itself as a variant of activation. For example, codeine (morphine 3-methyl ether) exhibits mainly antitussive and mildly analgesic action. When codeine undergoes demethylation in the organism, it converts to morphine, which is a strong analgetic.

**Deintoxication** resembles deactivation and is a defense reaction to the toxic effect of a drug. For example, phenol is a toxic compound, while phenol sulphate, which is a product of phenol conjugation in the organism, is nontoxic.

**Toxication** shows up as an enhanced side effect due to a drug administered into the organism. By mechanism, toxification resembles activation. Occasionally, toxification is produced by “lethal” molecules synthesized from the introduced compounds during their metabolism in the organism. The lethal synthesis with the involvement of a xenobiotic leads to a metabolic block and to the death of organism. For example, the administered fluoroacetate enters the Krebs cycle in tissues to produce a toxic product, fluorocitrate, which blocks aconitate hydratase and interrupts conversion steps in the Krebs cycle. Toxification effects are taken into account in the development of chemicals against rodents and other vermin.

### **Localization of drug metabolism in the organism.**

Depending on the site of conversion of biogenous preparations and xenobiotics in the organism, the drug metabolism is classified into cavitary (enteral), extracellular (humoral), and cellular, or tissue, types of metabolism.

The cavity, or enteral, drug metabolism is effected by hydrolytic enzymes supplied to the cavity of gastrointestinal tract. Hydrolysis of biogenous preparations occurs with the involvement of pancreatic and intestinal digestive enzymes. Xenobiotics whose molecules contain peptide, carboxyester, glycoside, amide and phosphamide bonds are also liable to hydrolysis. This process involves proteolytic and lipolytic enzymes as well as enzymes capable of hydrolyzing glycoside bonds. In addition, a large group of esterases (e.g. carboxyesterases and phosphatases) and phosphamidases (involved in hydrolysis of phosphamide bonds in drugs) are found in the intestine. Trypsin, while being a proteolytic, exhibits also an esterase activity and is capable of hydrolyzing the ester bonds in xenobiotics.

Extracellular, or humoral, drug metabolism takes place in the extracellular fluids (after uptake and subsequent circulation of a drug in the organism), i.e. in the blood, lymph, cerebrospinal, and extracellular proper, fluids. Possibly, metabolic conversions therein are chiefly confined to hydrolysis of the preparations delivered (both biogenous and xenogenous). In the blood and other fluids, this function is performed by proteinases and esterases (e.g. pseudocholeline esterase, phosphatases). In the extracellular fluids, other enzymes, for example, alcohol dehydrogenase, aminooxidases, etc., are available in small amounts, but the activity of these enzymes is rather low. The contribution of the humoral metabolic link to the overall drug metabolism is insignificant. At the humoral level, drug hydrolysis plays a role in drug inactivation; this metabolic link should be taken into account.

Cellular (tissue) drug metabolism. In the cells, the whole varieties of metabolic transformations, including those of xenobiotics, are being accomplished. However, the substances, before being subjected to the action of enzymic systems, should be transported from the site of their introduction to the cells and allowed to penetrate the intracellular space through the cell membrane. Xenobiotics are transported by the same mechanisms as

biogenous substances. In the blood plasma, they either become dissolved in the liquid medium, or adsorbed, mostly on albumin. In a dissolved or in a protein-bound state, xenobiotics are delivered to the cells (tissues). They gain access to the cells mostly by simple and facilitated diffusion; large molecules enter the cells by endocytosis. Xenobiotics synthetically derived from biogenous substances can be actively transported across the cell membranes using natural substance transport systems.

Not all the tissues and organs are equally active when they convert xenobiotics. The most actively engaged organ is liver which is in possession of enzymes that perform modification and conjugation of drugs. The other organs and tissues are less active in the metabolism of xenobiotics.

The metabolic conversion of xenobiotics occurs in various organelles of the liver cells. The most powerful metabolic system is found in endoplasmic reticulum (in microsomes). The microsomes are fragments of endoplasmic reticulum that are formed, for example, on trituration of a tissue sample and spontaneously close into small bladder-like structures (vesicles). Thus, with reference on its localization, the metabolism of xenobiotics is differentiated into microsomal and extramicrosomal. The extramicrosomal metabolism occurs in hyaloplasm, lysosomes, peroxisomes, and mitochondria.

**The enzymic reactions conducive to conversion of xenobiotics** may be divided into the following major groups: 1) oxidation-reduction reactions; 2) hydrolytic reactions 3) synthetic reactions, or conjugation reactions; 4) other reactions (isomerization, ring opening, etc., which are effected by isomerases and lyases).

#### **Microsomal oxidation of substances.**

In the microsomes, there are found enzymic chains for oxidation of substances. These chains are represented by two short electron-proton transfer chains built into the membranes of endoplasmic reticulum or into microsomal membranes. Microsomal oxidation is connected with these chains. One of these chains is a monooxygenase oxidation chain (in which the source of electrons and protons is reduced NADP), and the other is a reductase oxidation chain, with reduced NAD as a supplier of electrons and protons. The source of  $\text{NADPH}^+$  in the monooxygenase chain is the pentose phosphate cycle, and the source of  $\text{NADH}^+$  is glycolysis.

The microsomal  $\text{NADPH}^+$ -dependent monooxygenase chain is composed of flavoprotein (FP2), with FAD for a coenzyme, and cytochrome  $\text{P}_{450}$ . Flavoprotein exhibits a  $\text{NADPH}^+$ -dehydrogenase activity, FAD acting as an acceptor for two protons and two electrons. From flavoproteins, electrons are transported onto cytochrome  $\text{P}_{450}$ , and protons are lost into the environment (cytosol). Cytochrome  $\text{P}_{450}$  is the terminal self-oxidizable link of this chain. Like all the cytochromes, it belongs to hemoproteins. Its protein moiety is represented by a single polypeptide chain. The molecular mass of cytochrome  $\text{P}_{450}$  is about 50,000. The  $\text{P}_{450}$  is capable of complexing with carbon monoxide, CO. The light absorption maximum for these complexes is at 450 nm; hence the name for the given cytochrome.

Cytochrome  $\text{P}_{450}$  performs a dual function: it activates molecular oxygen by transferring electrons onto it, and uses the activated oxygen to oxidize substances R, with the concomitant formation of water. Consequently, one oxygen atom adds to the oxidizable substance (RO), and the other, by accepting two  $\text{H}^+$  ions from the medium, makes up water. The  $\text{NADH}^+$ -dependent reductase oxidation chain occurs not only in the microsomal membranes; it is also available in the outer mitochondrial membrane, in the nuclear membrane, and in the erythrocytic cell membranes. The reductase chain is thus included




among the most rapid reactions of biological oxidation, but its function in the cell remains still unclear. The self-oxidizable component of this chain capable of activating the oxygen has never been identified either; quite probable that this function is exercised by cytochrome P<sub>450</sub> itself. The NADPH<sup>+</sup>- and NADH<sup>+</sup>-dependent chains can exchange electrons. For example, the electrons from FP2 and cytochrome b5 may be transferred onto cytochrome P<sub>450</sub> to be used in the oxidation of substrates.

## MICROSOMAL OXIDATION

Dioxygenases:  $R + O_2 \rightarrow RO_2$

Monooxygenases:  $RH + O_2 + NADP2H \rightarrow ROH + H_2O + NADP$   
(hydroxylases)



### Conjugation of xenobiotics, its mechanism and role.


The conjugation stage, or synthetic stage, is essential for the formation of nontoxic and easily excretable drug metabolites. By their mechanism, the conjugation reactions are divided into two groups:

**Reactions of I type.** Initially, conjugating agents, i.e. biomolecules, are activated and then transferred onto xenobiotics to form conjugates. This type of conjugation reactions occurs in all tissues of the organism.

**Reactions of II type.** Initially, a xenobiotic is activated to be transferred onto a conjugating biomolecule to form a conjugate. This conjugation type is of rare occurrence and is only observed in liver and kidney.

Various groups for conjugation reactions of I and II types are distinguished, depending on the nature of a conjugating species involved. In the I type reactions, glucuronide, sulphate, acetyl, methyl, thiosulphate conjugations are to be noted, and in the II type, glycine and glutamine conjugations.


**Glucuronide conjugation.** UDP-glucuronic acid is the source for glucuronic acid residues in this process. Endogenous substances and xenobiotics are subject to glucuronide conjugation (known are glucuronides of bilirubin, steroid hormones, vitamin D, etc.). Xenobiotics can enter glucuronide conjugation if they possess or have acquired, during modification, a hydroxyl, carboxyl, and amino group (commonly, in the aromatic ring), or, at least, a SH-group. The conjugation reaction proceeds with the participation of UDP-glucuronosyltransferase by the scheme:



Among xenobiotics (drugs and poisons), susceptible to glucuronide conjugation are phenols, polyphenols, phenolic steroids, aromatic amino acids, and others.

**Sulphate conjugation.** Active form of a conjugating agent is 3'-phosphoadenosine-5'-phospho-sulphate (PAPS for short). PAPS, which may also be designated as PAP~SO<sub>3</sub>H, is a source of labile sulphate groups used in the conjugation of natural compounds and xenobiotics. The natural substances as subject to sulphate conjugation include endogenous


toxic products, e.g. indole, scatol, phenol as well as steroids, iodothyronines, tocopherols, and others. The sulphate conjugation reaction proceeds with the involvement of a special enzyme, sulpho-transferase, according to the scheme:



**Acetyl conjugation.** The source of labile acetyl groups in this variety of conjugation reactions is acetyl~S<sub>co</sub>A, which is produced by degradation of carbohydrates, triacylglycerides, and amino acids. Endogenous substances and xenobiotics containing a free NH<sub>2</sub> group may be acetylated.

N-acetylation is an essential biochemical reaction in the synthesis of monosaccharide derivatives (N-acetylglucosamine, N-acetylgalactosamine, and neuramic acid) that are further used in the synthesis of heteropolysaccharides. N-acetylation is also a route to neutralization of biogenous amines: serotonin, histamine, GABA, and others. N-acetylation of histones and nonhistonic chromatin proteins is an important regulatory mechanism of DNA transcription. For endogenous substances, the only case of O-acetylation has been reported, which is a reaction of acetylcholine formation.

Xenobiotics possessing a free NH<sub>2</sub> group (commonly, on the aromatic ring) are subject to acetylation. This reaction is effected by means of a special acetyltransferase called arylamine-N-acetyltransferase. This enzyme exhibits a low specificity to xenobiotics to be acetylated. The reaction proceeds by the scheme:




Among the xenobiotics susceptible to acetylation, sulphonylamides, isonicotinic acid hydrazides, and aniline derivatives can be mentioned; these preparations are widely used in medical practice.

**Methyl conjugation.** In this reaction methyl groups derived from the active form of methionine, S-adenosylmethionine, serve as a conjugating agent. S-adenosylmethionine is a participant in numerous reactions of methylation of endogenous compounds. It is also a methyl group donor for conjugation reactions of xenobiotics (RXH), which proceed with the involvement of methyltransferases, according to the scheme:



Xenobiotics containing an NH<sub>2</sub> group or heterocyclic nitrogen, as well as OH and SH groups, are subject to methylation by addition of methyl groups to N, O, and S atoms. Among the preparations used in therapy, liable to methylation are mono- and polyphenols, and heterocyclic compounds of pyridine, quinoline, isoquinoline, and thiouracil type.

**Thiosulphate conjugation.** This kind of conjugation is used in the enzymic detoxication of cyanides. The transfer of sulphur from thiosulphate onto a cyanide ion is catalyzed by a specific enzyme, thiosulphatesulphide transferase:




**Glycine conjugation.** This reaction belongs to type II conjugations, which require a prior activation of the substrate rather than of the conjugating agent. In principle, any carboxylic acid can serve as a conjugation substrate. The mechanism of glycine conjugation may be exemplified by the formation of hippuric acid. According to the mechanistic concept of type II conjugation reactions, the initial step is activation of benzoic acid with the involvement of arylacyl~S<sub>co</sub>A synthetase. Then benzoyl (or, in a wider sense, any activated substrate in the reactions of this type) is transferred onto the glycine amino group. This process is catalyzed by acyl-N-glycine transferase, which is specific to acylation of only glycine, barring other amino acids.

Similarly, glycine conjugates of other compounds are formed: aromatic acids (nicotinic), phenyl-substituted acetic acids (phenylacetic and hydratropic), steroid acids (cholic and deoxycholic).

**Glutamine conjugation** is a rare variety of conjugation, distinctly observable in patients with phenylketonuria. In normal humans, the glutamine conjugation of xenobiotics has never been reported.

To detoxify heavy metals, the liver contains **metallothioneins**, a group of cysteine-rich proteins with a high affinity for divalent metal ions such as  $\text{Cd}^{2+}$ ,  $\text{Cu}^{2+}$ ,  $\text{Hg}^{2+}$ , and  $\text{Zn}^{2+}$ . These metal ions also induce the formation of metallothioneins via a special metal-regulating element (MRE) in the gene's promoter.



**Figure 20.** Biotransformations (Koolman, 2005)

### Factors affecting drug metabolism.

Drug metabolism is affected by a variety of factors. These include genetic, age- and organ-specific, neuroendocrine, environmental factors, and the manner a drug has been administered. The rate at which a drug supplied to the organism is metabolized is dependent on the number of enzymes involved in modification and conjugation of the drug. In enzymopathies associated with defective enzymes that are involved in drug metabolism, a decrease in the drug metabolism rate is observed. Molecular diseases due to a defective UDP-glucuronosyltransferase are known. These molecular diseases are characterized by the disturbed glucuronide conjugation not only of bilirubin, but of other endogenous substrates and drugs too. For this reason, the prescription of sulphanilamides, salicylates, and phenol-derived preparations, which are metabolized by glucuronide conjugation, leads to aggravated symptoms of the disease; even normal doses of these drugs produce a negative effect.

The age is an important factor in drug metabolism. In neonates and infants, the enzymic apparatus of xenobiotic metabolism is poorly developed. As the young organism develops, the physiological enzymic deficiency disappears, while hereditary enzymopathies in adult humans persist.

Liver is the major organ responsible for drug metabolism. Environmental factors such as light, ambient temperature, radiation have been noted to influence the drug metabolism. The action of these factors is accomplished indirectly, via the neuroendocrine system. Over 200 preparations are known to be drug metabolism enzymes, primarily microsomal ones.

They include butadion (antiinflammatory), amidopyrine (analgesic), novocain (local anesthetic), ethanol, and others. Phenobarbital (soporific) acts as the most powerful inducer. It drastically enhances the synthesis of microsomal oxidation enzymes in the liver by affecting the genetic apparatus of the liver cells. Phenobarbital elicits the synthesis of UDP-glucuronosyltransferase and facilitates the conjugation stage in the metabolism of various materials.

Inducers for drug metabolism enzymes are biogenic preparations such as thiamine, riboflavin and their coenzymes, carnitine, pantothenic acid, androgens, and anabolic steroids; preparations of progesterone and estrogens inhibit these enzymes.

### Ethanol toxicity and its metabolism

The major site of ethanol degradation is the liver, although the stomach is also able to metabolize ethanol. Most of ethanol is initially oxidized by alcohol dehydrogenase to form acetaldehyde. A further oxidation, catalyzed by aldehyde dehydrogenase, leads to acetate. Acetate is then converted with the help of acetate-CoA ligase to form acetyl-CoA. In addition to cytoplasmic alcohol dehydrogenase, catalase and inducible “microsomal ethanol-oxidizing system” also contribute to a lesser extent to ethanol degradation. The rate of ethanol degradation in the liver is limited by alcohol dehydrogenase activity. The calorific value of ethanol is 29,4 kJ/g. Alcoholic drinks – particularly in alcoholics – can therefore represent a substantial proportion of dietary energy intake.






Figure 21. Ethanol metabolism (Koolman, 2005)

## Toxicity of ethanol.

Ethanol is rapidly distributed throughout the body. A large amount is taken up by muscles and brain. In the brain, ethanol is deposited in membranes and influences receptors for neurotransmitters. The effect of GABA is enhanced, while that of glutamate declines. High ratio of  $\text{NADH}/\text{NAD}^+$  facilitates the conversion of pyruvate into lactate, acetoacetate into  $\beta$ -hydroxybutyrate, biogenic amines into alcohols, the shifting acid-base balance to acidosis. Accumulation of lactate increases renal threshold for uric acid. The increase  $[\text{lactate}]/[\text{pyruvate}]$  ratio leads to gluconeogenesis inhibition and to hypoglycemia development. Ethanol inhibits the metabolism of some drugs in liver (for example, barbiturates), because it competes with them for  $\text{P}_{450}$ .



**Figure 22.** Blood ethanol level (Koolman, 2005)



**Figure 23.** Liver damage due to alcohol (Koolman, 2005)

High ethanol consumption over many years leads to liver damage. Ethanol-related high levels of  $\text{NADH}+\text{H}^+$  and acetyl-CoA in the liver lead to increased synthesis of neutral fats and cholesterol. However, since the export of the form VLDLs is reduced due to alcohol, storage of lipids occurs (fatty liver). The increase in the fat content of the liver (from less than 5% to more than 50% of the dry weight) is initially reversible. But in chronic alcoholism hepatocytes are increasingly replaced by connective tissue. When liver cirrhosis occurs the damage of the liver finally reaches an irreversible stage, characterized by progressive loss of liver functions. Acetaldehyde, which is about 15 times more toxic than alcohol, enters into condensation reactions with biogenic amines to form endogenous alkaloids. Acetaldehyde together with dopamine leads to the formation of salsolinol as well as tetrahydropapaveroline and  $\beta$ -carbolines. They bind with opiate receptors, causing hallucinations, stimulation of "enjoyment" centre. This explains the development of pathologic attraction to alcohol. Competitive inhibition of acetaldehyde dehydrogenase (for example, disulfiram) blocks the breakdown of acetaldehyde. The simultaneous intake of alcohol and disulfiram leads to increased acetaldehyde levels in the blood. This results in perspiration, tachycardia, nausea, vomiting and even severe circulatory failure.

## CONTROL QUESTIONS

1. Functions of the liver. The biological role of the liver in nitrogen metabolism and biosynthesis of specialized proteins.
2. The biological role of the liver in carbohydrate and lipid metabolism.
3. The biological role of the liver in metabolism of vitamins. Digestion, storage and excretion of different metabolites.
4. Hemoglobin metabolism, its breakdown. Bile formation.
5. Biochemistry of jaundice (hemolytic, hepatic, obstructive): causes, clinical symptoms, differential diagnostics.
6. Hereditary jaundice: Crigler-Nayar, Gilbert, Dabin-Johnson syndromes. Neonatal physiological jaundice.
7. Biotransformation of xenobiotics and endogenous toxins. Microsomal oxidation.
8. Ethanol toxicity and its metabolism.

## LIST OF INFORMATION SOURCES

1. Biochemistry: Manual for medical students or interns/ L.D. Popova, A.V. Polikarpova – Kharkiv: KNMU, 2011. – 539 p.
2. Gavriiliuc Ludmila A. Biochemistry: lectures for students of Medical Departments. – Chisinau, 2011. – 139 p.
3. Janson L.W., Tischler M. Medical Biochemistry: The Big Picture / The McGraw-Hill Companies, 2012. – 431 p.
4. Koolman J. et al. Color Atlas of Biochemistry (2th Ed.) – Thieme, 2005. – 476 p.
5. Murray R.K., Granner D.K., Mayes P.A., Rodwell V.W. Harper's illustrated Biochemistry. Twenty-sixth International Edition, 2003. – 702 p.

## CHAPTER 3. BIOCHEMISTRY OF KIDNEYS AND URINE.

### WATER-MINERAL METABOLISM.

#### **Functions of the kidneys. Filtration, secretion, reabsorption, excretion.**

#### **The mechanism of urine formation. Renal clearance.**

In the adult human organism the mass of the two kidneys is about 300g. The primary function of the kidney is to provide for a constancy of the internal medium of the organism.

Two zones are distinguished in the renal tissue:

- 1) the outer, or cortical, zone colored brown-red;
- 2) the inner, or medullary, zone colored lilac-red.

The basic functional unit of renal parenchyma is the nephron. In humans, the two kidneys number about 2 million nephrons.

In the nephron three major processes occur:

- 1) filtration at the glomerulus;
- 2) tubular reabsorption;
- 3) tubular secretion.

#### **Glomerular filtration.**

Glomerular filtration is a passive process. The total renal blood flow is about 1300 ml/min in adult human males. In health, the mean filtration rate is 125 ml/min. The filtration rate is determined by the filtration pressure:

$$FP = CP - (OP + Caps P),$$

where FP – filtration pressure, CP – capillary pressure, OP – oncotic pressure,

Caps P – intracapsular pressure.

In health FP ~ 30 mm Hg.

Capillary pressure within the kidney is dependent not so much on the arterial pressure as on the lumen ratio of the "afferent" and "efferent" glomerular arterioles. The efferent arteriole is narrower (by about 30% in diameter) than the afferent arteriole. The lumen ratio regulation for these arterioles is effected primarily by the kinin system.


The primary urine, practically devoid of proteins, is produced by ultrafiltration of the blood plasma into the lumen of Bowman's capsule. In health, proteins as colloid particles are incapable of penetrating the capsular cavity of the glomerulus through the capillary wall. Approximately 180 L of primary urine is produced. The pores in the glomerular basal membrane, which are made up of type IV collagen, have an effective mean diameter of 2,9 nm. This allows all plasma components with a molecular mass of up to about 15kDa to pass through membrane. At increasing masses, molecules are progressively held back; at masses greater than 65 kDa, they are completely unable to enter the primary urine. This applies to almost all plasma proteins – which in addition, being anions, are repelled by the negative charge in the basal membrane.

#### **Reabsorption and Secretion.**

Only 1% of the total fluid, filtrated in the glomerulus, is converted into urine. 99% of water, sodium chloride, hydrocarbonate ions, amino acids, 93% of potassium ions and 45% of urea are reabsorbed in the renal tubules. The cells of the proximal segment of the nephron reabsorb glucose, amino acids, vitamins and electrolytes, 6/7 of the fluid constitutive of the primary urine is also subject to reabsorption in the proximal tubules. In the tubule, organic substances (e.g., glucose, amino acids, lactate and ketone bodies) are recovered by secondary

active transport. There are several group-specific transport systems for resorbing amino acids, with which hereditary diseases can be associated (eg, cystinuria, glycinuria and Hartnup's disease).

Additional regulated reabsorption of water,  $\text{Na}^+$  and  $\text{Cl}^-$  occurs in the distal tubules. These processes are controlled by hormones (aldosterone, vasopressin). In the distal tubules potassium, ammonium, hydrogen ions may be secreted into the lumen of the nephron.



**Figure 24.** Urine formation (Koolman, 2005)

Renal clearance is used as a quantitative measure of renal function. It is defined as the plasma volume cleared of a given substance per unit of time. Inulin, a fructose polysaccharide with a mass of 6 kDa that is neither actively excreted nor reabsorbed but it freely filtered, has a clearance of 120 mL/min in healthy individuals.

Sodium ions penetrate from the tubular lumen into the cell by passive transport and they are transported from cells to the extracellular fluid by means of  $\text{Na}^+$ -pump. About 80% of the ATP energy in channel's cells is spend on the  $\text{Na}^+$ - pump.

The water uptake in the proximal segments is effected passively, assisted by the active absorption of sodium ions.

### **Functions of kidney.**

They include: the regulation of water and salt balance; the maintenance of acid-base balance; the maintenance of osmotic pressure; the removal of final products of metabolism; metabolic function; hormonal function.



Metabolic function. Specific features of renal tissue metabolism. Kidney uses at least 8-10% of the total oxygen consumed by the human organism. Concentrating urine and transporting it through membranes are processes that require large amounts of energy. The kidneys therefore have very high energy demands.

In the proximal tubule, the ATP needed is obtained from oxidative metabolism of fatty acids, ketone bodies, and several amino acids. To a lesser extent, lactate and glycerol are also used. In the distal tubule and Henle's loop, glucose is the main substrate for the energy metabolism. The endothelial cells in the proximal tubule are also capable of gluconeogenesis. The substrates for this are mainly carbon skeletons of amino acids. Their amino group is used as ammonia for buffering urine. Enzymes for peptide degradation and the amino acid metabolism occur in the kidneys at high levels of activity (e.g., amino acid oxidases, amine oxidases, glutaminase).

The first stage of creatine synthesis is performed in the renal and pancreatic tissues. Glycine amidinotransferase (or arginine-glycine transaminase) catalyzes this reaction. The observation of this enzyme in the blood may be linked either to a renal disease or to pancreonecrosis. Hydroxylation of 25-hydroxycholecalciferol occurs in kidney.

Hormonal function. The kidney plays an important role as an incretory (endocrine) organ. The juxtaglomerular cells, located in the region of the vascular pole of the glomerulus, produce rennin. Renin, through the agency (see water-salt metabolism) of angiotensin, exerts influence on the blood pressure of the whole organism, and on the production of aldosterone and ADH. The kidney also elaborates erythropoietin which stimulates the red blood cell production (erythropoiesis). Erythropoietin is a glycoprotein hormone. Its biosynthesis in the kidney is activated in a number of stress states – hypoxia, loss of blood, shock etc. The kidney produces prostaglandins, which are capable of influencing the responsiveness of the renal cells to the action of certain hormones.







Figure 25. Renal hormones (Koolman, 2005)

Role of kidney in maintenance of osmotic pressure is provided by means of rennin-angiotensin system. In addition to this, kidney is the target of aldosterone and ADH action. 3 main mechanisms:

- 1) Secretion of hydrogen ions; and formation and conversion of carbonic acid.
- 2) Sodium ions reabsorption and conversion of disubstituted phosphates to monosubstituted phosphates. In blood monosubstituted-to-disubstituted phosphate ratio is 1:4; in glomerular filtrate 9:1; in urine of the distal segment of the nephron 50:1.


$$\text{NaH}_2\text{PO}_4/\text{Na}_2\text{HPO}_4 = 1/4 \text{ (blood)} = 9/1 \text{ (glomerular filtrate)} = 50/1 \text{ (urine)}$$

- 3) Renal production of ammonia and its use instead of other cation for neutralization of acid equivalents and their urinary discharge.



## Inulin/creatinine clearance.

The measurement of the glomerular filtration rate (GFR) (volume of fluid filtered from the afferent arteriole into the renal corpuscle per unit time) allows clinicians to determine the health of the kidneys and to quantify any degree of kidney or renal failure. GFR is most accurately measured by injection of inulin into the bloodstream, a polysaccharide molecule from plants, which is neither reabsorbed nor secreted by the kidney. A more practical method of estimating GFR, though, is to measure blood creatinine, a molecule derived in muscle from the breakdown of creatine phosphate, a rapidly available source of adenosine triphosphate (ATP) energy for muscle and brain. Creatinine is exclusively excreted by the kidneys, completely filtered at the renal corpuscle, and only small amounts are secreted into the peritubular capillaries. The natural occurrence of steady levels of creatinine in the blood and urine offers an easy way to establish GFR and, therefore, kidney function. Varying mathematical formulas allow for correction of known variations depending on the patient's muscle mass, age, gender, race, and/or size. The normal range of GFR is (100-130) ml/min/1.73 m<sup>2</sup> but this varies in children and older adults. A GFR of over 60 ml/min/1.73 m<sup>2</sup> is usually sufficient for normal health, although high blood pressure, diabetes, and other chronic diseases can decrease kidney function and result in chronic kidney disease (CKD). CKD is enumerated in six stages, depending on the patient's GFR.



**Figure 29.** Overview of nephron and urine production (Janson. 2012)

## Physical and chemical properties of urine.

### Composition of urine under normal and pathological conditions.

#### Physico-chemical properties of urine.

**Daily urine** (diurnal diuresis): 1000-2000 ml (1500 ml). Pathologic state: <500 ml and >2000 ml. **Polyuria** (increased excretion of daily urine): after large fluid intake; after dietary intake of nutrients stimulating diuresis (eg. water melon and pumpkin).

**Pathologic polyuria:** in renal diseases (chronic nephritides and pyelonephritides), diabetes mellitus, diabetes insipidus (up to 15 liters).

**Oligouria** (diminished excretion of daily urine): in insufficient fluid intake, in febrile state, vomiting, toxicosis, acute nephritis. **Anuria** is complete or nearly complete suppression of urinary excretion: in severely affected renal parenchyma, ureteral obstruction.

In health, the urinary excretion is larger in the day time than at night. The diuretic day-to-night ratio varies from 4:1 to 3:1. In certain pathologic states (early stages of cardiac decompensation, cystopielitis) the passage of urine is larger at night than during the day. This state is known as **nycturia**.

**Color.** Normal urine is yellow. Colorless urine is under diabetes insipidus, excessive drinking, taking diuretics. **Red (“meat slops”) color** is caused by **hematuria**. Hematuria can be renal (as symptom of glomerulonephritis, trauma, nephrolytiasis) and extrarenal (under cystitis and urethritis). Red color of urine is also observed at porphyria and hemoglobinuria. **Orange color** is observed after taking several vitamins. **Red-violet** color occurs after beet eating. **Green color** is the sign of increasing putrefaction processes in intestine, after ingestion of rhubarb. **Color of dark beer** is examined under hepatitis. Blue color is observed after ingestion of methylene blue or as a sign of Hartnup disease. **Black urine** is present in patients with alkaptonuria.

**Odour.** Normal urine shows the characteristic smell or smell of “meat broth”. Ammonia odour occurs under hyperammoniemia. Aceton smell is observed in diseases which are accompanied by accumulation of ketone bodies. “Mouse or mould” smell shows phenylketonuria. “Maple syrup” or “beer yeasts” smell is the sign of leucinosia (maple syrup urine disease). “Cabbage” or “cat’s urine” smell occurs under tyrosinemia. The smell of rotted fish is present under trimethylglycinuria or dimethylaminuria. “Dirty socks” odour occurs under isovaleric aciduria.

**Transparency.** Normal urine is transparent. Muddy urine is observed at pyuria, hematuria, proteinuria, crystalluria, bacteriuria.

**Density.** The relative urine density in the adult human is liable to diurnal variation within a fairly wide range (from 1,002 to 1,035). Most commonly the urine density ranges from 1,017 to 1,020. Low density: in diabetes insipidus. High density: in acute nephritis, in diabetes mellitus.

**Isostenuria** – urine density is equal to that of the primary urine, or ultrafiltrate (~1,010). It is observed only in a severe renal insufficiency. The isostenuria indicates a disturbed concentration function of the kidney. This state has been recorded in chronic nephritis, “contracted kidney”.

**pH.** The urine usually has a slightly acidic pH value (pH 5,3-6,5). However, the pH value is strongly affected by metabolic status. After ingestion of large amount of plant food, it can increase to over 7. In meat rich diet it is acidic. The decrease of pH is observed in febrile states; diabetes mellitus; in starvation. The increase of pH is observed in cystitis, pyelitis, curative intake of alkaline mineral water.

## **Characterization of urine components in norm and in pathology.**

The urinary excretion of various materials reflects alterations in the processes that occur in the kidney and other tissues and organs of the organism. The daily volume of final urine amounts to 1.0-2.0 liters and the dry weight of final urine is about 60 g. Since the urine is a filtrate of blood plasma, it appears expedient to consider the urinary concentrations of various groups of biological materials from the standpoint of their occurrence in the blood plasma.

**Proteins.** In norm, the daily urinary excretion of proteins amounts to about 30 mg, which is not detectable by common laboratory techniques and routinely specified as “traces, or absence of urinary proteins”. Among the urinary proteins, enzymes are also present. The origin of normal urinary proteins is different.

**In pathology,** the urinary protein concentration may be increased; depending on the location of the damage, prevalent in the urine may be either plasmic proteins, or cellular proteins of the urinary tract. In inflammatory renal diseases (glomerulonephritides), the permeability of the basal membrane of nephron glomerulus increases; proteins are filtered in an amount above normal and fail to be reab-sorbed completely. Disturbances in the tubular protein reabsorption (nephroses) are conducive to a similar pathology. For this reason, in patients with glomerulonephritides and nephroses the urinary excretion of proteins may vary from 1 to 15-40 g per day. Nonetheless, even in such a contingency, the urinary proteins concentrations are small and can be detected only using special techniques. For example, in pancreatitis, an enhanced activity of  $\alpha$ -amylase and trypsin is observed both in blood and urine.

### **Nonproteinic nitrogenous urinary components.**

**Urea** is a major nitrogenous component of the urine. The normal excretion of urea is 333 to 583 mmol per day, which accounts for 60% to 80% of the overall urinary nitrogen. An increased urinary concentration of urea is observed in the states with pronounced catabolism of proteins and other nitrogenous components (starvation, burns, traumatism, atrophy of tissues, etc.). A decreased excretion of urea is observed in affected liver (urea-producing organ) and in impaired in the blood (this state is called azotemia).

**Uric acid.** Normally, the urinary excretion of uric acid is 2.35 to 5.90 mmol per day. Its increased urinary concentration is observed in a diet rich in nucleic acids or as produced by breakdown of cells and tissues, for example, leucocytes in patients with leucosis.

**Creatinine.** In norm, the urinary excretion of creatinine is 4.4 to 17.6 mmol per day; variations in creatinine concentration are dependent on muscular development. Physiological excretion of creatinine is normal only in children. In adult humans, creatinuria is a sign of pathology (e.g. muscular dystrophy).

**Amino acids.** In norm, the urinary excretion of amino acids is 0.29 to 5.35 mmol per day (as based on nitrogen). The urinary concentrations of glycine, histidine, and alanine are higher than those of other amino acids. In pathology (e.g. burns, diabetes mellitus, affected liver, and muscular dystrophy) hyperaminoaciduria may occur. Hereditary hyperaminoaciduria is associated with defective proteins-carriers for amino acids in the proximal renal tubules. In a disordered amino acid tissue metabolism, the urinary excretion of normally nonexcretable amino acid metabolites occurs (e.g. homogentisine acid, in alcaptonuria; phenylpyruvic acid in phenylketonuria).

**Ammonium salts.** In norm, the urinary excretion of ammonia as a component of ammonium salts (ammonium chloride) is 30-60 mmol per day. In pathology, an increased

urinary elimination of ammonium salts may be observed (in diseases accompanied by acidosis). A diminished excretion of ammonium salts occurs in diseases associated with alkalosis, in renal diseases due to affected distal tubules in which ammoniogenesis takes place.

**Hippuric acid.** The urinary excretion of hippuric acid is dependent solely on the amount of ingested vegetable food, since in the organism this acid endogenically is not produced. Commonly, the daily urine contains to 5.5 mmol of hippuric acid.

**Indican** (indoxyl sulphuric acid). Normal urine contains indican (in the form of potassium indoxyl sulphate) in trace amounts. In detectable quantities, indican appears in the urine on excessive alimentary intake of meat products; it also occurs as a byproduct of putrefactive processes in the intestine.

**Nitrogenous pigments.** Representative of these is stercobilinogen, a product of hemoprotein breakdown. Stercobilinogen is convertible to stercobilin and normally is excreted in the urine. In pathology, urinary excretions contain bile acids and a variety of bile pigments, for example, in affected liver and in toxicoses conducive to hemolysis.

#### **Nitrogen-free components of urine.**

**Glucose and other monosaccharides.** In norm, the daily urine contains a more 0.3-1.1 mmol of glucose. Such amounts escape detection by conventional analytical techniques; for this reason, glucose is not reckoned as a component of normal urine. However, in excessive dietary intake of carbohydrates, when the glucose concentration in blood attains a threshold value, i.e. of the order of 8.3-8.8 mmol/liter, alimentary glucosuria may develop in the organism. In pathology, glucosuria occurs due either to an increased blood glucose concentration, or to a defective carrier protein involved in glucose reabsorption in the renal proximal tubules. The former case is the most commonly encountered in the clinic, for example, in diabetes mellitus or in steroid diabetes. The latter case is the so-called renal diabetes. For example, the occurrence of fructose or pentose in the urine (renal fructosuria or renal pentosuria) is indicative of affected transport systems of the renal tubules.

**Lactic and pyruvic acids.** In norm, the daily urinary excretions of lactic and pyruvic acids amount to 1.1 and 0.11 mmol, respectively. An increased concentration of lactic acid in the urine is observed under intensive muscular work and in hypoxia. An increased urinary excretion of pyruvic acid occurs in diabetes mellitus and in B<sub>1</sub> hypovitaminosis.

**Ketone bodies.** In norm, the daily urine contains 20 to 50 mg of ketone bodies. At this level, they are not detectable by the analytical methods currently employed in the clinic. In pathology, increased concentrations of ketone bodies, i.e. a state called ketonuria, occur in diabetes mellitus, steroid diabetes, and starvation.

**Mineral salts.** In norm, the daily urine contains (in mmol): sodium, 174-222; potassium, 61-79; calcium, 4.02-4.99; inorganic phosphorus, 33. In pathology, an increase in urinary excretion of sodium and a decrease in excretion of potassium are observed in the adrenal hypofunction; the reverse situation occurs in hyperaldosteronemia and when mineralocorticoids and glucocorticoids are prescribed as drugs. A decreased urinary concentration of calcium and a distinct phosphaturia are observed when large doses of vitamin D and parathyrin are administered; a high urinary loss of calcium is characteristic of rickets and hypoparathyroidism.

## **The biological role of water. Osmotic pressure.**

### **Disorders of water metabolism (dehydration, hyperhydration).**

**Water Metabolism.** Structure features: Water molecule is dipole. Hydrogen bonds are formed between water molecules. This explains high boiling temperature and high thermal capacity of water.

#### **Biological role of water:**

- 1) Water is universal dissolvent. Water polarity provides good solubility of different substances and electrolyte ionization in water. Water is chemically inert, therefore substances dissolved in it keep their chemical and biological properties.
- 2) Water plays an important role in supporting unique structure and functions of cell organelles.
- 3) Water is obligatory component of biochemical processes. All the reactions in the organism are performed either in presence, or with participation of water.
- 4) Water performs transport function.
- 5) It participates in supporting osmotic pressure.
- 6) Water is an important thermoregulation factor.

**Water content in organism.** The total amount of body water is about 60-65% in adults, and 72% in newborns. The proportion of weight as water declines with age and with increased body fat content. By the age of 60 men have 51,5% of water and women 45,5% of it.

**Body water distribution.** Body water is distributed between 2 main compartments: **intracellular** and **extracellular**. Intracellular water comprises 70-72% of total body water of normal healthy adults. Extracellular water includes that of interstitial fluid and that of plasma, lymph, cartilage, etc. Major differences in composition between intracellular and extracellular fluids are the following:

- 1) Potassium is the principal cation within cells, whereas sodium predominates in extracellular fluid.
- 2) Because of many phosphorylated organic compounds are present within cells, phosphate is the primary intracellular anion, chloride replaces it in extracellular fluids.
- 3) Finally, the intracellular protein concentration is higher than that of blood plasma.

**Water balance.** Water is essential nutrient factor. A loss of 12-25% of water leads to death. In a normal healthy person, total body water volume remains remarkably constant, fluctuating less than 1% of body weight per day, and this constancy is maintained in spite of large variations in water intake. Daily water intake is 2,5-3 l. It depends on age, occupation, climate, diet, etc. Water need is higher in children, than in adults, namely: in children it is 100-150 g per kg of weight; in adults it is 30-50 g per kg.

#### **Water sources:**

- 1) Exogenic water is 85%. This is food and drinking water.
- 2) Endogenic water (15%). This is metabolic water.

The oxidation of 100 g of each carbohydrate, protein and fat yields 55.6; 41.3 and 107.1 g of water, respectively, but the total amount of metabolic water is quite small (200-400 ml per day) relative to that ingested in food or drink.

#### **Water losses:**

Water is required to replace fluid lost through the skin, lungs and gastrointestinal tract and to accompany renal excretion of urea, salts and other osmotically active substances. The amounts of these obligatory losses vary significantly with climate, activity level, state of health, and diet. Hot temperature, dry climate, vigorous physical activity and fever increase

water losses from the skin and lungs. Great amount of water is secreted into gastrointestinal tract with juices. Water secreted into the gastrointestinal tract is usually reabsorbed, but diarrhea and other intestinal diseases can result in very large water losses and organism dehydration. Total urine volume generally depends on water intake, but a minimum amount of water – an obligatory volume – is required to accompany the excretion of osmotically active substances especially urea, sodium chloride.

### **Regulation of water-salt metabolism.**

Regulation of water-salts metabolism come to supporting: constant osmotic pressure; constancy of total water volume in organism and its distribution between different fluid spaces; constancy of ionic composition; acid-base balance.

**Constancy of ionic composition** is provided by means of systems of active transport.

**Distribution of water** between fluid spaces of the organism is generally determined by physico-chemical mechanisms. This process is influenced by the following factors:

- 1) Osmotic pressure. Gradient of molar concentrations between fluid spaces of organism is motive force of water current between them. Water will be transferred to water space with greater molar concentration.
- 2) Oncotic pressure. Decreasing protein content in blood plasma leads to edema.
- 3) Hydrodynamic pressure in vessels (is created by heart work).
- 4) Permeability of cell membranes.
- 5) Active biological transport of ions.

**Regulation of total water volume constancy and osmotic pressure of blood** is performed by neurohumoral way. Osmotic pressure of extracellular fluid in greater extent depends on [NaCl], therefore base mechanism of osmotic pressure regulation is linked with the change of excretion rate of water, or sodium chloride. Regulation of extracellular fluid volume is performed by simultaneous change of excretion rate both water and sodium chloride. Supply of water to the organism depends on the thirst sense. Center of thirst is located in dorsal and central nucleus of hypothalamus. The water excretion by kidneys is regulated by neurohumoral way with participation of antidiuretic hormone.


**Antidiuretic hormone (vasopressin)** is synthesized in special neurons of the hypothalamus from which it is transported to the posterior pituitary and is secreted directly in blood. This is nonapeptide. Vasopressin stimulates the contraction of the muscular tissue of blood vessels (the vasopressory action). However, its major function is water balance control. It stimulates the reabsorption of water in the renal tubules through the adenylyl cyclase system and increasing hyaluronidase activity. Vasopressin secretion is stimulated by increasing osmotic pressure and by considerable decreasing volume of extracellular fluid. It should be noted, that the system of osmotic regulation functions in very limited range: a change of osmolality by 1% only leads to vasopressin secretion, which corrects this change. With the blood volumestem get opposite signals (for examples, the loss of blood under hyponatremia conditionse, it must be decreased about 7-15 % before similar response reaction is arisen. If both system get opposite signals (for examples, the loss of blood under hyponatremia conditions), a "volume" regulation prevails over osmotic regulation.

In pathology, for example, in atrophy of posterior pituitary, **diabetes insipidus** develops, a diseased state manifested by an excessive urinary water discharge. Vasopressin has important value for restoration of total volume of fluid in the organism. But increased reabsorption of water without sodium by vasopressin is little effective in restoration of extracellular fluid because 2/3 of reabsorbed water enters into intracellular space.



The regulation of sodium concentration is necessary to support the constancy of extracellular fluid volume. This regulation is performed by **aldosterone** and **sodium uretic factor**. In kidney aldosterone increases the sodium ions reabsorption (together with chloride) in distal tubules. This leads to a delay of sodium chloride in organism. In **hyperaldosteronism** the surplus delay of sodium chloride leads to increasing osmotic pressure. This is a signal to vasopressin secretion. Vasopressin enhances water reabsorption in the kidneys. Accumulation of sodium chloride and water is observed. Extracellular fluid volume is increased. Under supporting normal osmotic pressure, blood pressure is increased.

The primary regulators of aldosterone production by the glomerulosa cells are the **renin-angiotensin system** and potassium. Sodium, neural regulation, ACTH, adrenoglomerulotropin (isolated from pineal gland) are also involved.



**Figure 30.** RAAS (Janson. 2012)

Main mechanism of the regulation of the aldosterone secretion is reninangiotensin system. **Renin** is proteolytic enzyme, which is synthesized in juxtaglomerular cells of the renal afferent arterioli. They are sensitive to blood pressure change, to change of Na<sup>+</sup> and Cl<sup>-</sup> concentration in the renal tubular fluid. Any combination of factors, that decreases fluid volume or decreases NaCl concentration, stimulates rennin release. Renal sympathetic nerves that terminate in the juxtaglomerulas cells mediate the central nervous system effects on rennin release (through the β-adrenoreceptors). Renin is able to convert angiotensinogen (α<sub>2</sub>-globulin, produced by liver) into decapeptide angiotensin I. Angiotensin-converting enzyme, a glycoprotein, removes two carboxyl terminal amino acids from the decapeptide angiotensin I to form angiotensin II.

Various nonapeptide analogs of angiotensin I and other compounds act as competitive inhibitors of converting enzyme and are used for treating renin-dependent hypertension. These are referred to as angiotensin converting enzyme inhibitors. Converting enzyme also degrades bradikinin, a potent vasodilator; thus, this enzyme increases blood pressure in two distinct ways.

**Angiotensin II** increases blood pressure by causing vasoconstriction of the arteriole and is a very potent vasoactive substance. It is the potent stimulator of aldosterone production. It causes thirst. Angiotensin II inhibits renin release from the juxtaglomerular cells by feed-back mechanism.

**Sodium uretic factor** is synthesized by cells of auricle of the heart: 1) It stimulates excretion of sodium ions; 2) It shows vasodilatory effect; 3) It inhibits aldosterone synthesis; 4) It inhibits renin release.

**Supporting of pH constancy.** Acid-base balance is the relation between concentrations of hydrogen and hydroxyl ions in liquids of organism. Under normal conditions pH of blood is 7.35-7.45; pH of intracellular liquid is lower, than extracellular one, and pH value inside cells of different types may be different, but constant for the given type of cells. pH in different compartments of one type cell may be different. Difference of pH inside cells of various types and in various compartments of the given type of cells is explained by the features of metabolism, by mechanisms of active transport, by election permeability of membranes. Regulation of acid-base balance is achieved by physico-chemical (buffer systems) and physiological mechanisms (lungs, kidney).

### **Water and mineral metabolism disorders.**

Disturbances of water and mineral metabolism are distinguished into water and electrolyte imbalance (dishydrations) and disturbances of metabolism of separate minerals. Water imbalance is distinguished into:

#### 1. Dehydration:

- primary (hyperosmolar, pure water depletion);
- secondary (hypoosmolar, pure salt depletion);
- mixed (isoosmotic, water and salt depletion).

#### 2. Water intoxication (hyperhydration):

- total hyperhydration;
- intracellular hyperhydration;
- extracellular hyperhydration.

**Primary dehydration** (pure water depletion). Causes:

- inadequate water intake (coma, dysphagia etc.);
- in infants great amount of water practically without electrolytes may be lost through lungs in hyperventilation, fever, acidosis;
- excessive loss of water by kidney in diabetes insipidus.

Osmotic pressure in extracellular space is increased. Water flows from intracellular space to extracellular space. Intracellular dehydration develops. Symptoms: thirst, oligouria, hyperosmia, azotemia due to oligouria, hypernatremia.

**Secondary dehydration** (hypoosmolar). Causes: excessive sweating (mainly NaCl depletion); vomiting, diarrhea, duodenal fistules, cholera, etc.; Addison's disease; vigorous use of diuretics. From hypoosmolar extracellular space water flows to cells. This leads to intracellular edema. Symptoms: thirst is absent; dryness of skin; decreasing turgor of skin; headache; collapse.

**Mixed dehydration** (isoosmolar) occurs: in bleeding; peritonitis; exudates; in burns. The volume of both extracellular and intracellular fluids is decreased.

### **Water intoxication (hyperhydration).**

**Total hyperhydration** occurs due to excessive water intake or insufficient water excretion. Water is accumulated in all water spaces. Causes: severe stagnant cardiovascular insufficiency; hypersecretion of ADH following administration of anesthetics for surgery, administration of narcotic drugs or in stress (including any surgery); excess of aldosterone (Conn's syndrome); excessive parenteral administration of fluids. Symptoms: headache, nausea, depression.

**Intracellular hyperhydration.** Causes: infusion of hypotonic solutions; excessive drinking; insufficient excretion of fluid in nephropathies.

**Extracellular hyperhydration** (edema syndrome) due to the accumulation of water in interstitial fluid. Causes: reduction in colloid-osmotic pressure; increase of hydrostatic pressure; disturbances of functioning the heart; allergic and inflammatory processes.

**Disturbances of metabolism of separate minerals** can be **primary** and **secondary** ones. **Primary** disturbances are caused by deficiency or excess of any minerals in diet. Examples: endemic goiter (deficiency of I), fluorosis (excess of F).

**Secondary** disturbances of mineral metabolism may be caused:

- by insufficient amount of protein-carrier (for example, Addison-Biermer disease);
- by lack of apoenzyme (for example, insufficiency of sulfite oxidase – Mo containing enzyme – leads to mental retardation);
- by hormonal disbalance (for example, hypofunction of adrenal cortex – Addison's disease, hyperaldosteronism – Conn's syndrome).

## **The biological role of Na, K, Cl, Ca, Mg and P, disorders of their metabolism.**

### **The biological role of microelements. Dyselementoses.**

Most of minerals (sodium and potassium are notable exception) form salts and other compounds that are relatively insoluble, they are not readily absorbed, and most ingested minerals are excreted in feces. Mineral absorption often requires specific carrier proteins; the synthesis of these proteins serves as an important mechanism for control of mineral levels in the body. Transport and storage also require specific binding to carrier proteins. Excretion of most minerals is accomplished by kidneys, but many minerals are also secreted into the digestive juice and bile and are excreted with feces.

**Calcium.** The human body contains more calcium than any of other essential minerals – about 1-1.5kg g in 70-kg adults. Functions:

- 1) Plastic function. About 99% of the total amount of Ca is in bones and teeth. Most skeletal calcium is deposited as a form of hydroxyapatite,  $\text{Ca}_{10}(\text{PO}_4)_6(\text{OH})_2$ , but bone also contains considerable amounts of noncrystalline calcium phosphates and carbonates as well as a small amount of other salts. Because bone is constantly being remodeled, its mineral levels reflect the equilibrium between daily deposits and withdrawals. As much as 700 mg calcium may enter and leave the bones every day.
- 2) It is necessary for blood clotting.
- 3)  $\text{Ca}^{2+}$  takes part in coupling of excitation and contraction in muscle.
- 4) It provokes initial mediator secretion during synaptic excitation.
- 5) It is necessary for adequate functioning of membrane channels.
- 6) It is the "second messenger" that mediates cellular responses to a wide range of stimuli similar to the regulatory actions of cyclic nucleotides. The action of calcium appears to be mediated by intracellular receptor protein, calmodulin that binds calcium ions when their concentration increases in response to a stimulus. Calmodulin has been found to be present in every nucleated cell type examined. When  $\text{Ca}^{2+}$  is bound to calmodulin it modulates the activities of a great variety of enzymes, including those involved in cyclic nucleotide metabolism, protein phosphorylation, secretory function, muscle contraction, glycogen metabolism.

### **Phosphorus.**

- 1) This participates in bone formation (50% all the phosphorus in organism).

- 2) Phosphorus takes place in maintaining acid-base balance.
- 3) It is also an integral compound of nucleic acids, nucleotides, nucleotide coenzymes, phospholipids, some proteins, 2,3-bisphosphoglycerate.
- 4) It participates in formation of macroergic phosphate bonds (ATP, creatine phosphate etc.).

**Ca and P metabolism regulation.** Many hormones influence the calcium and phosphorus metabolism. Parathyroid hormone, calcitonin and hydroxylated forms of cholecalciferol are main of them. Normal plasma contains the equivalent of 9-11 mg of calcium per deciliter (2,25-2,75 mmol/L). The symptoms of calcium deficiency include tetany and related muscle and neurologic disorders. Low serum  $\text{Ca}^{2+}$  levels occur in vitamin D deficiency, hypoparathyroidism or renal insufficiency. The net negative  $\text{Ca}^{2+}$  balance leads to rickets in children or osteomalacia in adults. High serum  $\text{Ca}^{2+}$  levels accompany clinical disorders such as hyperparathyroidism, vitamin D intoxication, sarcoidosis, and cancer. Osteoporosis, which mainly occurs in women following the menopause, is based (at least in part) on a reduction in estrogen levels. Estrogens normally inhibit the stimulation of osteoblast differentiation by osteoblasts. If the effects of estrogen decline, osteoclasts predominate and excess bone removal occurs.

**Sodium.** Sodium is the cation ( $\text{Na}^+$ ) of the extracellular fluid. The concentration of  $\text{Na}^+$  in blood plasma is 126-152 mM, in erythrocytes 13,4-21,7mM. 1-3.5 g of Na is required daily for adults. Infants need 0.1-0.5 g and children 0.3-2.5 g daily. Functions of sodium:

- 1) Maintaining and supporting normal water balance and distribution of water in the organism.
- 2)  $\text{Na}^+$  is also important in the maintenance of osmotic pressure of body fluids and thus in protection against excessive fluid loss.
- 3) It is largely associated with chloride and bicarbonate in the regulation of acid-base equilibrium.
- 4) It participates in formation of resting membrane potential and regeneration of excitation potential.
- 5) It participates in supporting normal neuromuscular excitability.
- 6) It facilitates formation of conformation of enzyme molecule which is needed to precise orientation of catalytic groups.
- 7) It intensifies proteins swelling and increases the amount of water bound with them.

Every 24 hours approximately 2500 mmol of sodium are filtered by the kidneys. However, due to tubular reabsorption less than 1% of this sodium appears in the urine (100-200 mM/day). Approximately 80% of the filtered sodium are reabsorbed in the proximal tubules with equivalent amount of water. The reabsorption of sodium in the distal tubules is 5 times less than in the proximal ones. The reabsorption in the distal tubules takes place contrary to a concentration gradient and is regulated by aldosterone.

**Hyponatremia** may be relative and absolute. Relative hyponatremia is caused by excessive drinking or excessive parenteral infusion of hypotonic solutions. Absolute hyponatremia is caused by diuretic medication, intensive sweating, prolonged vomiting, intestinal fistula, in hypoaldosteronism.

**Hypernatremia** may also be relative and absolute. Relative hypernatremia is observed in hyperosmolar dehydration, in limited liquid intake. Absolute hypernatremia is developed in excessive Na intake, in primary hyperaldosteronism (Conn's syndrome), in secondary hyperaldosteronism, in chronic nephritis. The secondary hyperaldosteronism is caused by

hyperproduction of renin. This mechanism is observed in hypertonic disease, in cardiovascular insufficiency, etc.

**Potassium.** Potassium is the major intracellular cation. The concentration of potassium is 3.8-5.4 mM in blood serum and 79.8-99.3 mM in erythrocytes. Functions:

- 1) It participates in formation of resting membrane potential.
- 2) It favors the activation of some enzymes. It is needed for the synthesis of proteins, ATP, glycogen.
- 3) Potassium potentiates the function of parasympathetic nervous system and the action of acetylcholine on the nerve terminals in muscles.
- 4) It is involved in supporting acid-base balance.
- 5) It plays an important role in cardiac function.

**Hypokalemia** is observed in insufficient potassium intake, in changing redistribution of potassium between intracellular and extracellular fluids and in increased loss. Insulin facilitates the supply of potassium in cells, therefore hypokalemia is observed after infusion of insulin to patients with Diabetes mellitus. Loss of  $K^+$  in gastro-intestinal diseases, renal and extrarenal polyuria, diuretic therapy, primary and secondary hyperaldosteronism, metabolic and respiratory alkalosis causes hypokalemia. Hypokalemia is accompanied by muscle hypotonia, weakness, paresthesia, change of contractive function of myocardium, tachycardia. Potassium deficiency results in changes of electrochemical gradient of cell membranes of myocardium, a decrease of potential difference and depolarization of membrane. This leads to increasing muscular excitability. Low resting membrane potential causes slow enhancing and low amplitude of action potential. Therefore excitation irradiation is reduced. Myocardium insufficiency develops.

**Hyperkalemia.** The mechanism for excretion of potassium in normal persons is so effective that it is difficult to produce hyperkalemia simply increasing the oral intake. Hyperkalemia however may occur after rapid intravenous infusion of potassium salts. Hyperkalemia is also caused by excretion disturbance and by sudden release of potassium from the intracellular space. Decreased excretion of potassium is observed in renal failure, in hypofunction of adrenal cortex (Addison's disease). Damage of body cells from any cause results in release of cell contents including  $K^+$  into extracellular fluid. Crush injuries with damages to large volumes of muscle tissue, massive hemolysis are examples. In ketoacidosis there is substantial loss of intracellular  $K^+$  to the extracellular fluid. If ketoacidosis presents for a long time, there will be major depletion of total body  $K^+$ . In hyperkalemia the changes in myocardium are observed. Bradycardia, arrhythmia, blockade, asystole occur.

#### **Classifications of chemical elements, which are found in the organism:**

Macroelements: Na, K, Ca, Mg, P, S, Cl.

Microelements (trace elements): Fe, Cu, Zn, F, I, Se, Cr, Mn, Co, Mo.

Ultramicroelements: Br, Si, Sr, Li, Hg, Au, Ni, Ti, Ra, Cd, As, Ti, Sn etc.

Each element has to be supplied in organism in optimal concentration. Decreased supply results in diminishing biochemical process intensity. Increased supply leads to toxicosis.

## Summary of Important Minerals (Janson 2012)

Mineral	Function	Disease(s)
<b>Na</b>	Involved in maintenance of fluid volume and osmotic pressure per the kidneys and associated hormones (e.g., renin, aldosterone, antidiuretic hormone, atrial natriuretic peptide). Essential for generation and maintenance of electric or transport potential across membranes (e.g., nerve conduction, muscle contraction, and membrane pumps)	Hyponatremia. Neurological symptoms secondary to cell swelling and electrolyte imbalance; potentially fatal. Hypernatremia. Deficit in free water in the body. Variable symptoms, including neurological, potentially fatal.
<b>K</b>	Usually the partner to sodium, essential for generation and maintenance of electric and transport potential across membranes (e.g., nerve conduction, muscle contraction, and membrane pumps), as well as potassium-specific pumps.	Hypokalemia and hyperkalemia. Muscle and neurological symptoms; both may lead to fatal abnormal heart rhythm, especially hyperkalemia.
<b>Cl</b>	Involved in conjunction with sodium in maintenance of fluid volume and osmotic pressure per the kidney. Essential role in neurological functions (e.g., glycine and GABA neurotransmitters) and acid-base balance via transport of bicarbonate.	Hypochloremia and hyperchloremia. Often secondary to vomiting and/or diarrhea; usually asymptomatic but may have respiratory symptoms.
<b>Ca</b>	Required for bone formation and remodeling; important cofactor for several enzymes and signal for signaling pathways (i.e., diacylglycerol/ IP <sub>3</sub> ), including blood clotting and muscle contraction; neurotransmitter for some neuron signals and plays a prominent role in maintaining a potential difference across membranes	Hypocalcemia. Neurological symptoms; may be followed by potentially fatal spasms of larynx and abnormal heart rhythm. Hypercalcemia. Constipation (groans), psychotic episodes (moans), pain in bones, kidney stones, and depression, etc. (psychiatric overtones); abnormal heart rhythm can also develop.
<b>Mg</b>	Magnesium stabilizes phosphate groups, including those in ATP; cofactor in several enzymatic processes	Hypomagnesemia. Muscle weakness, nerve problems/ tremors, psychiatric episodes/ epileptic fits; may lead to heart failure. Hypermagnesemia. Weakness, breathing problems, and potentially fatal heart rhythms.
<b>P</b>	Essential structural and functional element for nucleic acids, bone/ teeth, and phospholipid component of membranes; addition or removal of phosphate to/ from a protein/enzyme serves as a key regulator of enzymes	Hypophosphatemia. Nerve, bone, red and white blood cells, membrane, and muscle functional problems. Hyperphosphatemia. (Interference with other minerals, promotes calcification of soft tissue organs).
<b>Fe</b>	Essential cofactor in numerous enzymes and proteins (e.g., heme); essential for oxidation processes or oxygen transport	Iron deficiency (anemia). Iron excess (hemachromatosis).
<b>I</b>	Essential element for thyroid hormones; can act as antioxidant outside of thyroid, may play a role in the development of breast and/or stomach cancer, and affects immune system and salivary gland health	Iodine deficiency (goiter; cretinism).

<b>Zn</b>	Cofactor in almost 100 enzymes, serving a multitude of roles in metabolism, transcription and translation, acid-base balance, immune function, and protein synthesis; part of unique, tertiary protein structures (e.g., zinc fingers); part of nerve response of glutamate and essential for learning	Zinc deficiency. Directly impacts the enzymatic processes that rely on it; initial signs may be seen in skin, hair, and nails. Zinc excess. Can impair the absorption of other ions (e.g., iron and copper); corrosive damage to soft tissues.
<b>Mn</b>	Essential cofactor for several types of enzymes involved in numerous biological functions as well as several specific types of peptides	Manganese deficiency. Possible association with inflammatory diseases, diabetes, and some neurological and psychiatric problems. Manganese excess (manganism). Progressive neurological/ psychiatric symptoms.
<b>Cu</b>	Cofactor in several enzymes involved in electron transport or oxidation-reduction reactions (e.g., cytochrome c oxidase). Also, used for electron transport	Copper deficiency (anemia symptoms, decreased metabolism, and psychiatric manifestations). Copper excess. Wilson's disease (neurological and psychological effects).
<b>S</b>	As part of cysteine and methionine amino acids, plays an essential role in component of primary and tertiary protein structure via disulfide bond as well as role in sulphur-containing enzymes (e.g., cytochrome c oxidase, coenzyme A (CoA)); reduction of reactive species via glutathione	NA
<b>Co</b>	Component of cobalt-containing cofactors/ enzymes, the most prominent of which is vitamin B <sub>12</sub>	Cobalt excess. Potentially fatal. Cobalt deficit (pernicious anemia).
<b>Ni</b>	Important cofactor in some enzymes (e.g., urease), especially those involved in reduction reactions	Nickel deficiency. Potential impact on involved enzymes, although not manifested as symptoms. Nickel excess. Skin irritant and potential cancer-causing agent.
<b>Cr</b>	Possible role in carbohydrate and/ or lipid metabolism	Chromium deficiency (extremely rare; effects controversial). Chromium excess (Cr <sup>3+</sup> . damage to DNA; Cr <sup>6+</sup> . can act as a cancer-causing agent and damages internal organs).
<b>F</b>	Role in strengthening of teeth and bone and, as such, used for prevention of cavities and treatment of osteoporosis	Fluoride deficiency. Possible connection to weakened teeth and bones. Fluoride excess. Neuromuscular and other symptoms that can result in death.
<b>Se</b>	Essential cofactor for certain antioxidant enzymes (e.g., glutathione peroxidase), which remove reactive oxygen species; believed to be cofactor in thyroid hormone conversion of T <sub>4</sub> to T <sub>3</sub>	Selenium deficiency. Rarely seen but may contribute to destruction of heart or connective tissue; also affects thyroid hormone synthesis Selenium excess (selenosis). Affects liver and lungs; potentially fatal.
<b>Mo</b>	Cofactor in several enzymes, including oxidizing enzymes (e.g., xanthine oxidase)	Affects enzymes requiring cofactor; neurological symptoms may result; possible association with development of esophageal cancer.

## CONTROL QUESTIONS

1. Functions of the kidneys. Filtration, secretion, reabsorption, excretion. The mechanism of urine formation. Renal clearance.
2. The role of the kidneys in the regulation of osmotic pressure and acid-base balance. Endocrine renal function.
3. Physical and chemical properties of urine. Composition of urine under normal and pathological conditions.
4. The biological role of water. The distribution of water and electrolytes in the body, its regulation. Osmotic pressure.
5. Disorders of water metabolism (dehydration, hyperhydration): types, causes, clinical symptoms.
6. The biological role of sodium, potassium and chlorine, regulation and disorders of their metabolism.
7. The biological role of calcium, magnesium and phosphorus, regulation and disorders of their metabolism.
8. The biological role of microelements (Fe, Cu, Zn, F, I, S, Se, Cr, Mn, Co, Mo). Dyselementoses: causes, clinical symptoms.

## LIST OF INFORMATION SOURCES

1. Biochemistry: Manual for medical students or interns/ L.D. Popova, A.V. Polikarpova – Kharkiv: KNMU, 2011. – 539 p.
2. Gavriiliuc Ludmila A. Biochemistry: lectures for students of Medical Departments. – Chisinau, 2011. – 139 p.
3. Janson L.W., Tischler M. Medical Biochemistry: The Big Picture / The McGraw-Hill Companies, 2012. – 431 p.
4. Koolman J. et al. Color Atlas of Biochemistry (2th Ed.) – Thieme, 2005. – 476 p.
5. Murray R.K., Granner D.K., Mayes P.A., Rodwell V.W. Harper's illustrated Biochemistry. Twenty-sixth International Edition, 2003. – 702 p.



## RECOMMENDED INFORMATION SOURCES

### Main literature

1. Biochemistry: Manual for medical students or interns/ L.D. Popova, A.V. Polikarpova – Kharkiv: KNMU, 2011. – 539 p.
2. Koolman J. et al. Color Atlas of Biochemistry (2th Ed.) – Thieme, 2005. – 476 p.
3. Lehninger A. Principles of Biochemistry. – Sixth Edition. David L. Nelson and Michael M. Cox. – 2013. – 1336 p.
4. Rostoka L.M. «Medicine. Biological chemistry – Krok 1» Self-Study guide for students for licensing examination Krok 1 (medical care) / L.M. Rostoka, A.D. Sitkar, Ya.Yu. Burmistrova, H.E. Reyti – Uzhhorod, 2019. – 172 p.
5. Satyanarayana U., Chakrapani U. “Biochemistry”, Fourth Edition. – 2013. – 809 p.
6. USMLE™. Step 1 Biochemistry and Medical Genetics Lecture Notes – 2013 Kaplan, Inc. – 430 p.

### Optional literature

1. Berg J.M. et al. Biochemistry (5th Ed.) – Frreman, 2002. – 1515 p.
2. Biological chemistry. Lecture notes / A. D. Tahanovich [et al.]. – Minsk : BSMU, 2017. – 144 p.
3. Gavriliuc Ludmila A. Biochemistry: lectures for students of Medical Departments. – Chisinau, 2011. – 139 p.
4. Litwack G. Human biochemistry and disease (1th Ed.) – 2008. – 1272 p.
5. Mardashko O.O., Yasinenko N.Y. Biochemistry. Texts of lectures. – Odessa. The Odessa State Medical University, 2003. – 416 p.
6. Mary K. Campbell – Biochemistry (6th Edition) – 2007. – 836 p.
7. Murray R.K., Granner D.K., Mayes P.A., Rodwell V.W. Harper’s illustrated Biochemistry. Twenty-sixth International Edition, 2003. – 702 p.
8. Richard A Harvey, Ph. D.; Denise R Ferrier, Lippincott's Illustrated Reviews: Biochemistry, Fifth Edition. – 2011. – 531 p.
9. Janson L.W., Tischler M. Medical Biochemistry: The Big Picture / The McGraw-Hill Companies, 2012. – 431 p.
10. Lelevich S.V. Clinical biochemistry: manual for students of the Faculty of Foreign Students with English medium of instruction / S.V. Lelevich, T.V. Popechits. – Grodno: GrSMU, 2010. – 86 p.

### Internet sources

1. <http://e-learn.uzhnu.edu.ua>

# CONTENTS

FOREWORD.....	3
CHAPTER 1. BIOCHEMISTRY AND PATHOBIOCHEMISTRY OF BLOOD	
Blood as a specialized tissue of the body, its composition. Functions of blood.....	4
Physical and chemical properties of blood. Inorganic components of blood.....	5
Buffer blood systems. Acid-base balance, its regulation. Acidosis and alkalosis.....	7
Blood plasma proteins. Pathoproteinemia. Acute phase proteins. Lipoproteins. Non-protein organic compounds of blood. Azotemia.....	10
Blood plasma enzymes. Enzyme diagnostics.....	17
Respiratory function of erythrocytes. Transport of O <sub>2</sub> and CO <sub>2</sub> . Hemoglobin, its synthesis. Metabolism of porphyrins. Metabolism of iron. Disorders of hemoglobin metabolism: hemoglobinopathy, thalassemia, porphyria.....	21
CHAPTER 2. BIOCHEMISTRY OF LIVER. XENOBIOTICS AND DETOXIFICATION PROCESSES.	
The role of the liver in protein, carbohydrate and lipid metabolism. Biosynthesis of specialized proteins. Digestion, storage and excretion of different metabolites.....	32
Hemoglobin metabolism, its breakdown. Bile formation.....	38
Biochemistry of jaundice (hemolytic, hepatic, obstructive): causes, clinical symptoms, differential diagnostics.....	41
Biotransformation of xenobiotics and endogenous toxins. Microsomal oxidation.....	44
Ethanol toxicity and its metabolism.....	52
CHAPTER 3. BIOCHEMISTRY OF KIDNEYS AND URINE. WATER-MINERAL METABOLISM.	
Functions of the kidneys. Filtration, secretion, reabsorption, excretion. The mechanism of urine formation. Renal clearance.....	55
Physical and chemical properties of urine. Composition of urine under normal and pathological conditions.....	66
The biological role of water. Osmotic pressure. Disorders of water metabolism (dehydration, hyperhydration).....	63
The biological role of Na, K, Cl, Ca, Mg and P, disorders of their metabolism. The biological role of microelements. Dyselementoses.....	67
RECOMMENDED INFORMATION SOURCES.....	73