

574.58

RICCIOCARPUS NATANS (L.)CORDA (RICCIACEAE)

Ricciocarpus natans (L.)Corda (Ricciaceae)

Ricciocarpus natans.

: *Ricciocarpus natans*,
kunik@mail.Uzhgorod.ua

The rare communities with *Ricciocarpus natans* (L.) Corda (Ricciaceae) on the Transcarpathians lowland. – Felbaba-Klushyna L.M. – The rare communities with *Ricciocarpus natans* are describe on the Transcarpathians lowland. The characteristic of the phytocoenoses and the ecological conditions are consider.

Key words: *Ricciocarpus natans*, Transcarpathians lowland, water communities, reclamation canal.

Address: Uzhgorod National University, 32, A. Voloshyn St., Uzhgorog, 88018, Ukraine; e-mail: kunik@mail.Uzhgorod.ua

30% [2].
1860
50- [2].
Ricciocarpus natans
6
0,3-0,8
10-20 ².
Ricciocarpus natans
+ *Lemna minor*.
80 %, *Ricciocarpus natans* – 80%,
Lemna minor – 15% , *Lemna trisulca* – 5%,
Utricularia vulgaris – 2-3%, *Myriophyllum verticillatum* – 8%.
Carex riparia (40%), *Sparganium erectum* (20%), *Lycopus europaeus* (2-3%), *Bidens frondosa* (1%) *Typha latifolia*, *Glyceria maxima* (1%)
78 [1].
Spirodella polyrrhisa (1%).
Ricciocarpus natans.

(.) - [1],
 4-5 0,5-1 , - , , , -
Ricciocarpus -
natans + *Elodea canadensis*: *Ricciocarpus natans* , 40-100 , -
 50-60%, *Elodea* - -
canadensis - 15%, *Lemna minor* - 5-7%, *Lemna*
trisulca - 2%, *Utricularia vulgaris* - 5-7%, 19 , -
Myriophyllum verticillatum - 5%, *Spirodella* - 17 . -
polyrrhiza (3%). - , -
Typha latifolia. -
Sium latifolium *Polygonum amphibium*
Ricciocarpum natantis Slavni 1956, - var. *terrestris*.
Lemnion minoris, *Lemnetalia minoris*
 K. Tx. 1955, Lemneta R. Tx. 1955.

1. Ltmneta, Potametea, 2.
 Ruppiaetea, Zosteretea, Izoeto-Littorelletea, Phragmito- : . . . , 1964. - . 59-60.
 Magnocaricetea / // -
 . - : , 2006. - . 69-71.

: 10 2007 .
 : 27 2007 .

(*Aconitum nanum* (Baumg.) Simonk) -
 (Aconitum nanum (Baumg.) Simonk) 2007 -
 1350-1400
 (1991).
 (, , , 2003). - . . .