Vydavateľ / Publishing a /and adresa redakcie / Address of editor: VPS - SLOVAKIA, spol. s r.o., Lúčna 1A, 080 06 Prešov, Slovenská republika, IČO: 36457256 IČ DPH: SK2020011697 Obch. reg. Okr. súdu PO, vložka 11334/P

Číslo účtu / Account number: 2627729574 Kód banky 1100 Tatra banka, a.s. pobočka Prešov

IBAN: SK84 1100 0000 0026 2772 9574 Swiftový kód: TATRSKBX

Variabilný symbol /Variable symbol VS 114800 Konštantný symbol / Constant symbol KS 0308

Generálny partner / General Partner: MINISTRY OF EDUCATION AND SCIENCE OF UKRAINE STATE UNIVERSITY "UZHOROD NATIONAL UNIVERSITY" 46 Pidhirna St, Uzhorod, Transcarpathia, Ukraine, 88000

 \implies + 380312233341 \implies + 380312234202 \implies official@uyhnu.edu.ua Identification code 02070832 http://www.uzhnu.edu.ua/

Vedecká rada / Scientific board:

Predseda: Editor in Chief:	Tajomnik: Scientific	Podpredsedníčka: Vice
Dr.Sc Prof. Smolanka	Secretary:	Chairman:
Volodymyr	Dr.Sc Prof. Kostenko Yevhen	Dr.Sc Prof. Oksana Klitynska

Členovia: Scientific boards:

Dr. Fejérdy Pál, D.M.D., Ph.D Department of Prosthodontics	Dr.h.c., C.Sc.,Prof.Andrej Jenca MUDR(Kosice, Slovak
Faculty of Dentistry Semmelweis University (SE) (Budapest,	Republic)
Hungary)	
Dr.Sc.Prof.Slabkiy Genadiy (Uzhhorod, Ukraine)	Dr.Sc.Prof.Myronyuk Ivan (Uzhhorod, Ukrain)
Dr.Sc.Prof.Fera Olexandr (Uzhhorod, Ukrain)	Dr.Sc Prof. Rusyn Vasil (Uzhhorod, Ukraine)
Dr.h.c Prof. Hanna Eliasova, PhD (Presov, Slovak Republik)	Dr.Sc Prof. Rusyn Andryy (Uzhhorod, Ukraine)
Dr.h.c., Dzupa Peter, PhD, MUDR (Chadca, Slovak Republik)	Dr.Sc Prof. Boldizhar Patricia (Uzhhorod, Ukraine)
Dr.Sc. Prof. Savichuk Natalia(Kyiv, Ukraine)	Dr.Sc Prof. Boldizhar Oleksandr (Uzhhorod, Ukraine)
Dr.Sc Prof. Kasakova Rimma (Uzhhorod, Ukraine)	Dr.Sc Prof. Korsak Vyacheslav (Uzhhorod, Ukraine)
Dr.Sc Prof. Potapchuk Anatolyy (Uzhhorod, Ukraine)	Dr.Sc Doc. Rumyancev Kostyantin (Uzhhorod, Ukraine)
Dr.Sc. Prof. Tore Solheim (Oslo,Norway)	Dr.Sc. Prof. Bobrov Nikita (Kosice, Slovak Republic)
Dr.Sc Prof. Mishalov Volodymyr (Kyiv, Ukraine)	Dr.Sc. Prof. Hokan Mornsted (Sweden)
Dr. Sc. Prof. Vilma Pinchi (Florence, Italy)	Dr.h.c., Prof. Jozef Zivcak, PhD (Kosice, Slovak Republik)
Dr.Sc. Prof. Tkachenko Pavlo (Poltava, Ukraine)	Dr.Sc. Doc. Gasuk Petro (Ternopil, Ukraine)
Dr.Sc. Prof. Kaskova Liudmyla (Poltava, Ukraine)	Dr.Sc. Doc. Gasuk Natalia (Ternopil, Ukraine)

Vedecký časopis je registrovaný na Ministerstve kultúry Slovenskej republiky a УКРАЇНА Держана реєстраційна служба України N º 20221 – 10021Р

The Scientific journal registered at the Ministry of culture of the Slovak Republic.

Vedecký časopis. Kategória publikačnej činnosti **ADF** podľa prílohy č. 1 Smernice č. 13/2008-R zo 16. októbra2008 o bibliografickej registrácii a kategorizácii publikačnej činnosti, umeleckej činnosti a ohlasov Ministerstva školstva Slovenskej republiky. Vedecké práce v domácich nekarentovaných časopisoch. Články alebo štúdie, ktoré zverejňujú originálne (pôvodné) výsledky vlastnej práce autora alebo autorského kolektívu uverejnené v nekarentovaných vedeckých časopisoch.

The Scientific journal. A category of publishing activities pursuant to annex 1 of Directive no **ADF** No 13/2008-R from 16. October 2008 on the bibliographic entry registration and categorisation of the publishing activities of the Ministry of education of the Slovak Republic, artistic activity and responses. Scientific work in domestic not current journals. Articles or studies, which published the original work of the author or copyright (original) the results of its own staff members published in the not current scientific journals.

Periodicita: 4x ročne. Periodicity: 4x per year. Dátum vydania: August 2015. The date of issue: August 2015.

	TABLE OF CONTENT	
1	КРИТЕРІЇ ПОСТАНОВКИ ДІАГНОЗУ ТА ВИБІР МЕТОДУ ЛІКУВАННЯ ХРОНІЧНОГО ФІБРОЗНОГО ПУЛЬПІТУ ТИМЧАСОВИХ ЗУБІВ Клітинська О.В., Зорівчак Т.І., Гасюк Н.В.	4
2	10 ПРИЧИН ОБРАТИ КОМПОМЕРНИЙ ПЛОМБУВАЛЬНИЙ МАТЕРІАЛ НА ДИТЯЧОМУ СТОМАТОЛОГІЧНОМУ ТЕРАПЕВТИЧНОМУ ПРИЙОМІ Васько А.А., Шеверя С.,М., Іваськевич В.З.	9
3	АНАЛІЗ МЕТОДІВ ДОСЛІДЖЕННЯ КАРІЄСРЕЗИСТЕНТНОСТІ ЕМАЛІ ЗУБІВ У ДІТЕЙ Клітинська О.В., Шетеля В.В.	13
4	USAGE EFFECTIVENES OF IMPROVED ENDODONTO-ENDOOSSAL IMPLANTS FOR STABILIZATION OF FRONTAL TEETH Maistuk P., Pruts H., Mochalov Yu.	16
5	ПРАВОВЕ ЗАБЕЗПЕЧЕННЯ МЕДИЧНОЇ ДОПОМОГИ ДИТЯЧОМУ НАСЕЛЕННЮ В УКРАЇНІ Слабкий Г.О., Дудник С.В.	19
6	АЛГОРИТМ ПІДГОТОВКИ ЗАКЛАДІВ ОХОРОНИ ЗДОРОВ'Я ДО ЗМІНИ ПРАВОВОГО СТАТУСУ НА НЕПРИБУТКОВІ НЕКОМЕРЦІЙНІ ПІДПРИЄМСТВА Скрип В.В., Слабкий Г.О., Миронюк І.С.	24
7	ДО ПИТАННЯ ПРО БЕЗПЕКУ ПАЦІЄНТІВ В АНЕСТЕЗІОЛОГІЇ: ІНГАЛЯЦІЙНА АНЕСТЕЗІЯ СЕВОФЛУРАНОМ В УМОВАХ НЕДОСТУПНОСТІАНАЛІЗАТОРА КОНЦЕНТРАЦІЇ АНЕСТЕЗІОЛОГІЧНИХ ГАЗІВ Федосюк Р. М., Андерс А.В., Літвін Т.Д., Мельник А.В., Щупачинська Л.О., Щупачинський В.Б.	30
8	ІНТЕРАКТИВНІ МЕТОДИ У ВИКЛАДАННІ ЗАГАЛЬНОЇ ХІРУРГІЇ Філіп С.С., Скрипинець Ю.П.	41
9	FEATURES OF DIGITAL VISUALIZATION OF SCARRING TISSUE AND INDIVIDUAL-PERSONAL CHARACTERISTICS OF PATIENTS WITH HEAD AND NECK HYPERTROPHIC SCARS Bukhanchenko O.P., Avetikov D.S., Ivanyts'ky I.O.	45
10	THE DYNAMICS OF CHANGES OF INDICATORS OF PRODUCTS OF PEROXIDE LIPID OXIDATION IN THE GOMOGENATES OF SCARRING TISSUES IN DIFFERENT TERMS OF THE POSTOPERATIVE PERIOD Krynychko L.R., Grygorov S.M., Stavickiy S.O., Lokes K.P., Rozkolupa O.O.	50
11	МІКРОФЛОРА ПОРОЖНИНИ РОТА У ХВОРИХ НА ГЕНЕРАЛІЗОВАНИЙ ПАРОДОНТИТ І СТУПЕНЯ ТЯЖКОСТІ Савєльєва Н.М., Соколова І.І.	55
12	ФОРМУВАННЯ МАНУАЛЬНИХ ГІГІЄНІЧНИХ НАВИЧОК У ДІТЕЙ ДОШКІЛЬНОГО ВІКУ М. УЖГОРОДА. Мельник В.С., Горзов Л.Ф., Гриненко Є.М., Горзов В.В.	61
13	МОЖЛИВОСТІ МОДИФІКАЦІЇ ПІДХОДУ ДО ЛІКУВАННЯ ПАРОДОНТИТУ ВИХОДЯЧИ ІЗ ПРИНЦИПІВ Р-4 МЕДИЦИНИ Костенко Є.Я., Савчук О.В., Бунь Ю.М.	67

UDC: 351.77:616.314-053.2+577.118

USAGE EFFECTIVENES OF IMPROVED ENDODONTO-ENDOOSSAL IMPLANTS FOR STABILIZATION OF FRONTAL TEETH

Maistuk P.*, Pruts H.**, Mochalov Yu.***

*- assistant Professor of Department of Pediatric Dentistry, Uzhhorod National University, Ukraine. ** – Doctor Clinic of Maxillo-Facial surgery, Comenius University in Bratislava, Slovakia

***- Candidate of Medical Science, Associate Professor of the Department of Surgical Dentistry, Maxillo-Facial Surgery and Oncology, Faculty of Dentistry, Uzhhorod National University, Uzhgorod, Ukraina

Summary: Despite the rapid development of materials and treatment methods for caries and its complications (pulpitis, periodontitis), its prevalence among the Ukrainian population remains periodontitis, which is accompanied by the destruction of the alveolar bone part, leads to the need for surgical treatment of the disease (resection of the tooth root apex) for the preservation of teeth, especially the frontal group. Known ways to improve the biological parameters of teeth after surgical resection of the root top are their reinforcement with the help of endodonto-endoossal implants. This study shows the evaluation of the effectiveness of surgical treatment with the use of advanced endodonto-endoossal implants among young people who live in the Transcarpathian region. Data analysis of the evaluation of the effectiveness of the proposed method of surgical treatment of patients has made it possible to establish its high efficiency and recommend it as a method of choice.

Key words: endodonto-endoossal implant (EEI), periodontitis, people of young age, Transcarpathian region, effectiveness evaluation, root apex resection (RAR), bone density, HU (Hounsfield Unit), Periotest, conventional units (CU).

Actuality. Despite the rapid development of methods of treatment and rehabilitation of patients in dentistry, the prevalence of caries and its complications, among the Ukrainian population, remains at a high level [1, 7, 12]. That is why endodontic treatment of teeth occupies a significant place among all therapeutic dental interventions [2, 8, 9,]. According to the official reports of the Ministry of Health of Ukraine, the level of dental morbidity among the Ukrainian population has increased over the last ten years, which is especially noticeable among children and young people and leads to complications. It is this fact that causes the improvement of the techniques of radical interventions to restore the structure and functioning of the tooth-jaw system [6, 11, of the common surgical 13]. interventions is RAR, which is used in the treatment of the frontal group of teeth, which also have an aesthetic function as an element of the quality of life of the patient [4], especially among young people.

However, teeth after resection of the root top the biochemical parameters significantly reduce, increases the functional load, movement of the tooth is found, and often this factors leads to its loss [3, 10].

A known method for improving the biomechanical parameters of stabilization following surgical root top resection interventions is their reinforcement by endodonto-endoossal implants [5, 14], especially when it comes to the frontal group of teeth among young people, where the use of implants after resection of the tops is an alternative method treatment with preservation of aesthetic and functional result in different periods of postoperative term.

The aim of the study. Evaluation of the effectiveness of the surgical treatment performed with the use of advanced endodonto-endoossal implants.

Results and discussion. In order to determine the effectiveness of the proposed method for restoring the biological function of the frontal group teeth after resection of the root top using an advanced endodonto-endossal implants, a group of 73 patients aging 16-24 years old with the following

diagnoses were selected: chronic granulomatous periodontitis of the frontal upper teeth group and the lower jaw (the size of the defect of the bone tissue at the apex of the root is up to 0.3 cm), chronic granulomatous periodontitis of the teeth frontal group of the teeth of the upper and lower jaws (size of the defect in bone on the top of the root is from 0.3 cm to 0.5 cm). Depending on the choice of treatment method, all patients were divided into 4 clinical groups. For all clinical groups of patients, additional methods of examination were performed.

Determination of the degree of mobility of the established EEI was carried out using the technique "Periotest" using the apparatus ("Periotest-S") produced by "Siemens" (Germany) [15]. The evaluation of the results were carried out on the basis of the received digital data using the Miller scale in the modification of Flesrar (1980).

Computed tomography was used to determine the bone density, and the evaluation of the data was performed using the X-ray absorption scale (the Hounsfield scale) and was ultimately expressed in Hounsfield units [15]. The evaluation of efficacy in all clinical groups was performed in several stages - 7 days after surgery and in long-term follow-up, in particular after 3, 6 and 12 months after treatment. Density of bone tissue was recorded during the initial examination, 7 days after the treatment (surgery), after 3, 6 and 12 months after the treatment. The average mobility indicators in 3 months indicate a reliable lower mobility of teeth in patients of the third clinical group, where as an alternative treatment method, an improved method of RAR with additional filling of bone tissue defect with an osteoplastic material was chosen (+ 29.0 ± 1.26 C.U., 18.0 ± 0.21 C.U., $+ 10.0 \pm 0.8$ C.U.,, $+ 24.0 \pm 1.24$ C.U., p <0.05). These figures correspond to the first and fourth groups - 2 degree of mobility, in the second -1-2 degree, and in the third group 0-1 degree of mobility.

In the first and fourth control clinical groups, after 6 and 12 months of follow-up, no significant dental stabilization and a

decrease in the degree of mobility were found, which remained at the level of 1-2 degree. The results obtained in the first clinical group ((+ 27.0 \pm 1.24 and + 24.0 \pm 1.24 conventional units and the fourth control group (+ 24.0 \pm 0.2 and + 20.0 \pm 1.31 conventional units)) confirm this assertion.

In the second and third clinical groups there is a stable positive dynamics with a clear tendency to reduce the degree of mobility of teeth (the second clinical group with indicators + 16.0 \pm 0.21 and + 10.9 \pm 0.11 conventional units) to 0-1 degree. In the third clinical group, after 12 months, it was almost possible to reduce the mobility of the teeth after surgery for 0 degree (+ 4,0 \pm 0,01 and + 2,0 \pm 0,01 conventional units, respectively).

The degree of mobility of teeth is confirmed by the results of densitometry. Restoration of the density of bone tissue at the site of defect to norm was observed among patients of the second and third groups reliably in 12 months. After 3 and 6 months after the treatment in the control and first clinical groups, the results in the dynamics were different (346.4 ± 1.13 HU and 310.2 ± 1.34 HU in the control and 540.0 \pm 2.33 HU and 620.0 \pm 2,38 HU - in the first clinical groups). After 12 months, the results in both clinical groups did not differ significantly (769.0 \pm 1.35 HU; 743.0 \pm 3.34 HU; p> 0.05), however, they significantly differed from those of healthy jawbone tissues (p < 0.05).

In the second clinical group, after 3 and 6 months after surgery, the density of bone tissue in the defect area was 756.3 ± 2.45 HU and 880.1 ± 4.34 HU. In the third clinical group, these indices were insignificantly higher and amounted to 834.3 ± 4.35 HU and 880.4 ± 2.56 HU. 12 months after the treatment, in the second clinical group, the density of bone tissue was 957.4 ± 2.45 HU, and in the third group, 1003.3 ± 4.38 HU.

Thus, the use of EEI of the improved design in a complex with the filling of bone defect by osteoplastic material after 6-12 months after conducted RAR restores the density of bone tissue in the defect zone to the

indicators of the density of healthy bone tissue of the jaw (880,1 \pm 4,34 HU; 1003,3 \pm 4,38 HU; 1000 \pm 2,45 HU; p> 0,05).

In the course of statistical calculations it was established that the index of bone density in the zone of defect in the first and control groups increases unevenly, however, 12 months after treatment reached the maximum relative to all previous results of the markings.

When constructing Euclidean distances diagrams, it was found that for 12 months at initially close bone density in the defect area, there was a significant

improvement in all groups of patients (p <0.05), although the effectiveness and uniformity of bone resection differed.

Conclussion. The statistical relationship between the mobility of the teeth and the density of bone tissue in the defect zone can be interpreted as a direct dependence. This testifies to the effectiveness of the proposed methodology and the feasibility of its use in practical surgical dentistry and to serve as a method of choice.

REFERENCES

- 1. Біденко Н. В. Патогенез, клінічна картина, прогнозування, особливості лікування і профілактики карієсу зубів у дітей віком до 3 років // автореф. дис. на здобуття наук. ступеня д. мед. наук: спец. 14.01.22 стоматологія. Київ. 2012. 35 с.
- 2. Бланкова С. Л. Протокол ведения больных с хроническим генерализованным пародонтитом / С.Л. Бланкова, Н.А. Макарова // Практическая медицина (стоматология). Журнал для практикующих врачей. Казань, 2011. № 1 (33). С. 63—67.
- 3. Галахин К.А. Применение материала Bio—Oss при костно-пластических операциях / К. А. Галахин, П. В. Сидельников // Современная стоматология. 2013 №4. C.101–102
- 4. Загорский В.А. Биомеханика костной ткани челюстей и черепа человека / В.А. Загорский, В.В. Загорский // ДенталЮг. Краснодар, 2010. № 5. С. 64—67.
- 5. Клітинська О.В. Комплексне обгрунтування ранньої діагностики, профілактики та поетапного лікування карієсу у дітей, які постійно проживають в умовах біогеохімічного дефіциту фтору та йоду/ Дисертація на здобуття наукового ступеня доктора медичних наук. Ужгород 2015. 337с.
- 6. Клітинська О.В. Епідеміологічні аспекти поширеності основних стоматологічних захворювань дітей Закарпаття / Клітинська О.В. // Інноваційні технології в стоматології: мат. наук.-практ. конф. (м. Тернопіль, 9–10 жовтня 2012 р). Тернопіль: ТДМУ ім. Горбачовського, 2012. С. 34
- 7. Клітинська О.В. Ранжування уражень карієсом зубів у дітей при ретроспективному аналізі первинної медичної документації/ О.В.Клітинська, А.А. Васько// Україна. Здоров'я нації. 2016. № 3 (39). С. 39-44.
 - 8. Стоматологічна допомога в Україні / під ред. Князевич В.М. К.,2009. 34 с.
- 9. Хоменко Л.О. Терапевтична стоматологія дитячого віку / Л.О.Хоменко, О.І.Остапко, О.Ф.Кононович, В.І.Шматко та ін. Підручник.— К.: Книга плюс, 2013. 766 с
- 10. Черникис А.С. Особенности хирургических методов введения эндооссальных, субпериостальных эндодонт-эндооссальных конструкций металлических имплантатов в целях ортопедического лечения в стоматологии: дис. на соискание научной степ. кандидата мед. наук: спец. 14.01.14 стоматология./ Черникис А.С. М., 1988. 173 с.
- 11. Gapski R. Effect of systemic matrix metalloproteinase inhibition on periodontal wound repair: a proof of concept trial / R. Gapski, J.L. Ban, D.P. Sarment // J. Periodontol. 2014. Vol. 75, № 3. P. 493–494.
- 12. Klitinska O.V. Efficiency estimation of using phased program of caries prevention in children domiciled in Transcarpathian region / O.V.Klitinska, Y.Ya. Kostenko, Y.A. Mukhina, A.A. Vasko, N.V. Layosh// Acta stomatologica Naissi. − 2016. − Vol. 32, № 74. − P. 1635-1649.
- 13. Klitinska O.V. Exogenus drug prevention of dental caries in primery school children with high caries activity / O.V.Klitinska, A.A. Vasko, Y.A. Mukhina, // Intermedical journal. 2016. Vol. I (7). P. 10-12
- 14. Loe H. Oral hygiene in the prevention of caries and periodontal disease / Loe H. // Int dent j. 2010. Vol. 50. P. 129–139.
- 15. Ultrastructural characterization of the implant/bone interface of immediately loaded dental implants / Meyer U, Joos U, Mythili J, [at al.] // Biomaterials. 2014. 25 P. 1959–1967.