



Wiadomości Lekarskie

Official journal of the Polish Medical Association



Memory of
dr Władysław
Biegański

VOLUME LXXV, ISSUE 5 PART 2, MAY 2022

Since 1928



ALUNA Publishing House



Wiadomości Lekarskie

Editor in-Chief:

Prof. Władysław Pierzchała

Deputy Editor in-Chief:

Prof. Aleksander Sieroń

Statistical Editor:

Dr Lesia Rudenko

Managing Editor:

Agnieszka Rosa – amarosa@wp.pl

International Editorial Office:

Nina Radchenko (editor)

– n.radchenko@wydawnictwo-aluna.pl

Polish Medical Association (Polskie Towarzystwo Lekarskie):

Prof. Waldemar Kostewicz – President PTL

Prof. Jerzy Woy-Wojciechowski – Honorary President PTL

International Editorial Board – in-Chief:

Marek Rudnicki

Chicago, USA

International Editorial Board – Members:

Kris Bankiewicz	San Francisco, USA	George Krol	New York, USA
Christopher Bara	Hannover, Germany	Krzysztof Łabuzek	Katowice, Poland
Krzysztof Bielecki	Warsaw, Poland	Jerzy Robert Ładny	Białystok, Poland
Zana Bumbuliene	Vilnius, Lithuania	Henryk Majchrzak	Katowice, Poland
Ryszarda Chazan	Warsaw, Poland	Ewa Małecka-Tendera	Katowice, Poland
Stanislav Czudek	Ostrava, Czech Republic	Stella Nowicki	Memphis, USA
Jacek Dubiel	Cracow, Poland	Alfred Patyk	Gottingen, Germany
Zbigniew Gasior	Katowice, Poland	Palmira Petrova	Yakutsk, Russia
Mowafaq Muhammad Ghareeb	Baghdad, Iraq	Krystyna Pierzchała	Katowice, Poland
Andrzej Gładysz	Wrocław, Poland	Tadeusz Płusa	Warsaw, Poland
Nataliya Gutorova	Kharkiv, Ukraine	Waldemar Priebe	Houston, USA
Marek Hartleb	Katowice, Poland	Maria Siemionow	Chicago, USA
Roman Jaeschke	Hamilton, Canada	Vladyslav Smiiianov	Sumy, Ukraine
Andrzej Jakubowiak	Chicago, USA	Tomasz Szczepański	Katowice, Poland
Oleksandr Katrushov	Poltava, Ukraine	Andrzej Witek	Katowice, Poland
Peter Konturek	Saalfeld, Germany	Zbigniew Wszolek	Jacksonville, USA
Jerzy Korewicki	Warsaw, Poland	Vyacheslav Zhdan	Poltava, Ukraine
Jan Kotarski	Lublin, Poland	Jan Zejda	Katowice, Poland

Distribution and Subscriptions:

Bartosz Guterman prenumerata@wydawnictwo-aluna.pl

Graphic design / production:

Grzegorz Sztank

www.red-studio.eu

Publisher:

ALUNA Publishing House

ul. Przesmyckiego 29,

05-510 Konstancin – Jeziorna

www.wydawnictwo-aluna.pl

www.wiadomoscilekarskie.pl

www.wiadlek.pl

CONTENTS

ORIGINAL ARTICLES

- Olga Hadław-Klimaszewska, Agnieszka Jankowska, Justyna Laskowska, Marta Woldańska-Okońska
USING THE SODA SCALE TO ASSESS THE EFFECTIVENESS OF NEUROLOGICAL SPEECH AND LANGUAGE THERAPY ON IMPROVING LANGUAGE FUNCTIONS IN POST-STROKE PATIENTS 1229
- Aidyn G. Salmanov, Alla D. Vitiuk, Orusia A. Kovalyshyn, Serhiy M. Baksheev, Tetiana V. Kutytyska, Svitlana M. Korniyenko, Victor O. Rud
PREVALENCE AND RISK FACTORS OF INFERTILITY IN UKRAINE: RESULTS A MULTICENTER STUDY (2019-2021) 1234
- Israa H. Alfuaadi, Ibrahim A. Altamemi
IMPACT OF FGFR4 (GLY388ARG) GENE POLYMORPHISM ALONG WITH VISFATIN CYTOKINE AND HIGH MOBILITY GROUP BOX-1 (HMGB1) ON ACUTE CHOLECYSTITIS 1242
- Volodymyr B. Fik, Ruzhena M. Matkivska, Yosyp M. Fedechko, Vasyl V. Humeniuk, Oksana V. Yefremova, Larysa Ya. Fedoniuk
INTERDEPENDENCE OF THE MICROBIOTIC COMPOSITION OF BIOPELLICLE AND THE SEVERITY DEGREE OF CHANGES IN THE MUCOSA OF THE GUMS AFTER TEN WEEKS OF EXPERIMENTAL OPIOID EXPOSURE 1248
- Ibtisam H. Al-Azawi, Mahasin S. Al-Bidiri
DISTRIBUTION OF INTEGRON III AND PHYLOGENIC CLADE AMONG MDR UROPATHOGENIC *E. COLI* FROM PATIENT IN AL-DIWANIYAH CITY, IRAQ 1254
- Irina Pinchuk, Vitaliy Pishel, Marina Polyvianaiia, Oksana Kopchak, Stanislav Chumak, Natalia Filimonova, Yuliaya Yachnik
EMOTIONAL IMPACT OF THE COVID-19 PANDEMIC ON HEALTHCARE WORKERS IN UKRAINE 1261
- Ansam Zeki Thaker, Lubna Amer Al-Anbari, Essraa Mohsen Al-Essawe
DUAL TRIGGER STRATEGY AFTER CONTROLLED STIMULATION INTRAUTERINE INSEMINATION CYCLE DID NOT INFLUENCE THE PREGNANCY OUTCOME COMPARED WITH STANDARD HCG TRIGGER ONLY PROTOCOL 1268
- Vladyslav A. Smiiianov, Tetiana V. Fartushok, Lesia A. Rudenko, Nadiia V. Fartushok
THE ROLE OF LEPTINRESISTANCE IN THE PATHOGENESIS OF OBESITY IN PREGNANT WOMEN 1274
- Sally B. Shabeeb, Shaima R. Ibraheem, Kareem M. Lilo, Yaqoob A. Saleh, Noor A. Kazim
EVALUATION OF SOME BIOCHEMICAL MARKERS AND THE ANTIBODY RESPONSE TO HEPATITIS B VACCINE IN HEMODIALYSIS PATIENTS 1284
- Kateryna Pivtorak, Iryna Fedzhaga, Natalya Pivtorak, Larysa Vozniuk, Olexandra Klekot
FAT AND MUSCLE COMPONENTS OF BODY WEIGHT AND THEIR RELATIONSHIP WITH THE CONCENTRATION OF SERUM ADIPOKINES IN PATIENTS WITH NONALCOHOLIC FATTY LIVER DISEASE 1289
- Hussein A. Al-Khairallah, Mohammed H. AL-Yasiri
MOLECULAR DETECTION OF THE TWO VIRULENCE GENES HWP1 AND ALS1 IN CANDIDA SPECIES ISOLATED FROM ONYCHOMYCOSIS 1295
- Oleksandr S. Avramchuk, Oksana Yu. Plevachuk, Iryna A. Koval
BEREAVEMENT AND COVID-19: PREVALENCE, COMORBIDITY, AND ASSOCIATED FEATURES AMONG UKRAINIAN SAMPLE 1299
- Ali Faris Abdul Hussein, Alaa Khalaf Awad, Burhan Hadi
NURSES' KNOWLEDGE ABOUT INFECTION CONTROL AT PRIMARY HEALTH CARE CENTERS IN AL-HILLA CITY, IRAQ 1305
- Volodymyr Hryn, Yuriy Kostylenko, Natalia Svintsytska, Valentyna Bilash, Volodymyr Lytovka
THE ISSUE OF HISTOLOGICAL IDENTIFICATION OF M-CELLS IN THE PEYER'S PATCHES OF ALBINO RAT SMALL INTESTINE 1309
- Shurooq Wesam Al-Shaibani, Hayfaa Jaber Hussein, Hala Kadhim Jawad, Waleed J. A. Al-Kelaby, Sanaa Abdul-Razzaq Ibrahim Al-Rubaie
PHYSIOLOGICAL AND HISTOLOGICAL STUDY OF THE CALCIUM OXIDE NANOPARTICLES EFFECT ON THE TESTIS OF MALE WISTER RATS 1313
- Olexandr V. Shumakov, Olena V. Dovhan, Tetiana V. Talaeva, Iryna V. Tretiak, Olexandr M. Parkhomenko, Olga S. Gurjeva
THE PROGNOSTIC UTILITY OF LEUKOCYTE AND PLATELET COUNTS FOR RISK ASSESSMENT OF IN-HOSPITAL COMPLICATIONS IN PATIENTS WITH ACUTE ST-SEGMENT ELEVATION MYOCARDIAL INFARCTION 1317
- Veronika M. Dudnyk, Iryna I. Andrikevych, Katerina Khromykh, Halyna I. Mantak, Henadii M. Rudenko
ANTIBIOTIC ASSOCIATIVE DISORDERS OF THE MICROBIOTIC OF THE COLON IN INFANTS WITH ACUTE RESPIRATORY DISEASES 1323
- Oleksandr Pulyk, Myroslava Hyryavets, Taras Studeniak
POSTSTROKE FATIGUE AND MOTOR RECOVERY AFTER ISCHEMIC STROKE 1328

ORIGINAL ARTICLE

POSTSTROKE FATIGUE AND MOTOR RECOVERY AFTER ISCHEMIC STROKE

DOI: 10.36740/WLek202205218

Oleksandr Pulyk, Myroslava Hyryavets, Taras Studeniak

UZHGOROD NATIONAL UNIVERSITY, UZHGOROD, UKRAINE

ABSTRACT

The aim: To determine the effect of post-stroke fatigue on the motor recovery after ischemic stroke.

Materials and methods: The study was conducted on the basis of the department of vascular neurology of the Uzhhorod Central City Clinical Hospital during the six months of 2020. We examined 92 patients who suffered a hemispheric ischemic stroke in the early recovery period.

Results: At the end of therapy, improvement of motor function was observed in patients in both groups. We received a significant improvement ($p < 0,05$) in the patients of the main group on the SSS and 6-point scale of muscle strength. We found that a high score on the FAS showed a significant correlation with the worst dynamics of recovery on the NIHSS, Scandinavian scale of stroke and 6-point scale of muscle strength ($p < 0,05$).

Conclusions: According to the results of the study, it was found that the presence of post-stroke fatigue significantly impairs the improvement of motor functions in patients with hemispheric ischemic stroke, patients who had more severe fatigue had worse recovery rates after stroke.

KEY WORDS: ischemic stroke, paresis, post-stroke fatigue

Wiad Lek. 2022;75(5 p2):1328-1330

INTRODUCTION

Poststroke fatigue is a common symptom in patients. A systematic review and meta-analysis found marked variabilities in estimates of poststroke-fatigue prevalence, ranging from 25% to 85%[1]. Poststroke fatigue is generally qualitatively different from fatigue experienced before stroke, as the former can be exacerbated by stress and physical exercise, and generally responds well to rest and adequate sleep[2,3]. Prospective study has reported that, at 6 months after stroke, approximately half (51%) of all patients complained of fatigue[4]. Regarding the duration of fatigue after stroke, acute fatigue can last up to 6 months, whereas the chronic type can persist in 40% of patients after 2 years[2]. Several studies have reported poststroke fatigue to be an independent predictor of shorter survival, institutionalization, poorer functional outcome, and greater dependency for activities of daily living along with instrumental activities of daily living[2]. Moreover, in young patients, poststroke fatigue has been reported to be a determinant to resuming work, independent of physical disability or cognitive deficit[5]. In addition, in patients aged 18 to 50 years, poststroke fatigue has been associated with a poor functional outcome, as assessed by the modified Rankin Scale (odds ratio, 4.0; 95% CI, 1.6–9.6)[6].

THE AIM

The aim was to determine the effect of post-stroke fatigue on the motor recovery after ischemic stroke.

MATERIALS AND METHODS

The study was conducted on the basis of the department of vascular neurology of the Uzhhorod Central City Clinical Hospital during the six months of 2020. We examined 92 patients who suffered a hemispheric ischemic stroke in the early recovery period. The criteria for inclusion in the study were: age from 40 to 74 years, verified diagnosis of ischemic stroke of hemisphere localization, absence of severe somatic diseases, poststroke fatigue (the average score on the FAS more than 22 points). General clinical examination, neuroimaging examination (computed tomography, magnetic resonance imaging of the brain), neuropsychological examination using a battery of tests: the Scandinavian scale of stroke (SSS), NIHSS, Scale for assessing muscle strength (Peak L. M. And Braddom R, 1996), the Ashworth scale for assessing the tone of muscles and the Fatigue Assessment Scale (FAS) were used.

Rehabilitation was carried out according to the protocol in the early recovery period during the month with the use of physical therapy, occupational therapy, constraint-induced movement therapy, physiotherapy treatments.

Statistical calculations are performed using the standard statistical package "Microsoft Excel 2010". Differences at $p < 0,05$ (95% significance level) and at $p < 0,01$ (99% significance level) were considered statistically significant.

RESULTS

We examined 92 patients who suffered a hemispheric ischemic stroke in the early recovery period (the first 3




Fig. 1. Dynamics of the functional state of patients in Scandinavian scale stroke

Table I. The effect of poststroke fatigue on motor recovery after ischemic stroke

Scales	FAS	
	the main group	the control group
Scandinavian scale of stroke	r=0,51, p=0,0018	r=0,48, p=0,0016
NIHSS	r=0,38, p=0,0095	r=0,43, p=0,0018
6-point scale of muscle strength, hand	r=-0,43, p=0,0017	r=0,39, p=0,0014
6-point scale of muscle strength, leg	r=0,34, p=0,005	r=0,38, p=0,004
Ashworth's scale, hand	r=0,13, p=0,22	r=0,22, p=0,13
Ashworth's scale, leg	r=-0,10, p=0,49	r=0,21, p=0,14

months). The average age of patients was $59,1 \pm 1,1$ years. Analyzing the location of brain damage, it was found that 49 patients (54%) had an ischemic stroke of the left hemisphere, and 43 patients (46%) - right hemisphere. The patients were randomly divided into two groups of 46 patients. In the main group physical therapy, occupational therapy, constraint-induced movement therapy, physiotherapy treatments were used, in the control group we used the same methods, except constraint-induced movement therapy.

Before the start of rehabilitation therapy, the mean NIHSS score in patients of the first group was $13,3 \pm 2,8$ points, in patients of the control group - $12,5 \pm 3,7$ points. The Scandinavian scale of stroke in patients in the main group had an average score of $42,6 \pm 0,8$ at baseline and $44,2 \pm 0,7$ points in the control group (fig. 1).

According to the 6-point scale of muscle strength, for the hand the average score for patients in the first group was $2,9 \pm 0,7$ points, for patients in the control group $3,0 \pm 0,8$ points. According to Ashworth's scale, the muscle tone in the paretic limb of the patients in the main group before the study was $2,4 \pm 0,8$ points, and in the control group patients - $2,2 \pm 0,6$ points. According to the the Fatigue Assessment Scale average score for patients in the first

group was $29,1 \pm 3,7$ points, in the control group - $28,2 \pm 2,9$ points.

At the end of therapy, improvement of motor function was observed in patients in both groups. We received a significant improvement ($p < 0,05$) in the patients of the main group on the SSS and 6-point scale of muscle strength.

We found that a high score on the FAS showed a significant correlation with the worst dynamics of recovery on the NIHSS, Scandinavian scale of stroke and 6-point scale of muscle strength ($p < 0,05$) (Table I)

DISCUSSION

The current study was aimed to determine the effect of post-stroke fatigue on the motor recovery after ischemic stroke. Poststroke fatigue is a common symptom that can have debilitating effects. Fatigue is defined as a feeling of lack of energy, weariness, and aversion to effort [1]. Fatigue is common after stroke with a prevalence ranging from 16% to 72% [1-2] depending on the population studied and whether patients with mood disorders are included or excluded. Moreover, up to 40% of stroke survivors report it as their worst or one of their worst symptoms [1-2]. The

chronic type can persist in 40% of patients after 2 years [2]. Our study was conducted in patients who suffered a hemispheric ischemic stroke in the early recovery period during the six months of 2020. To detect fatigue in patients we used the FAS – a scale to assess post-stroke fatigue. This scale is the most frequently used scale in patient with stroke[3]. Approximately half (51%) of all patients complained of fatigue at 6 months after stroke[4]. Poststroke fatigue to be an independent predictor of poorer functional outcome and greater dependency for activities of daily living. In our study according to the the Fatigue Assessment Scale average score for patients in the first group was $29,1 \pm 3.7$ points, in the control group - $28,2 \pm 2.9$ points. To detect disorders of the nervous system we used the Scandinavian scale of stroke(SSS), NIHSS, Scale for assessing muscle strength (Peak L. M. And Braddom R, 1996), the Ashworth scale. Before the start of rehabilitation therapy, the mean NIHSS score in patients of the first group was 13.3 ± 2.8 points, in patients of the control group - 12.5 ± 3.7 points. According to the 6-point scale of muscle strength, for the hand the average score for patients in the first group was 2.9 ± 0.7 points, for patients in the control group 3.0 ± 0.8 points. Rehabilitation was carried out according to the protocol in the early recovery period during the month with the use of physical therapy, occupational therapy, constraint-induced movement therapy, physiotherapy treatments. At the end of therapy, improvement of motor function was observed in patients in both groups. We received a significant improvement($p < 0,05$) in the patients of the main group on the SSS and 6-point scale of muscle strength. A number of studies suggest that in patients poststroke fatigue to be a determinant to resuming work, independent of physical disability or cognitive deficit [2-7]. Our study also showed that the patients who had more severe fatigue had worse recovery rates after stroke.

CONCLUSIONS

According to the results of the study, it was found that the presence of post-stroke fatigue significantly impairs the improvement of motor functions in patients with hemispheric ischemic stroke, patients who had more severe fatigue had worse recovery rates after stroke.

REFERENCES

1. Cumming T.B., Yeo A.B., Marquez J. et al. Investigating post-stroke fatigue: an individual participant data meta-analysis. *J Psychosom Res.* 2018;113:107–112. doi: 10.1016/j.jpsychores.2018.08.006.

2. Paciaroni M., Acciarresi M. Poststroke Fatigue. *Stroke.* 2019;50(7):1927–1933. doi:10.1161/STROKEAHA.119.023552.
3. Ozyemisci-Taskiran O., Batur E.B., Yuksel S. et al. Validity and reliability of fatigue severity scale in stroke. *Top Stroke Rehabil.* 2019;26:122–127.
4. Hawkins L., Lincoln N.B., Sprigg N. et al. The Nottingham Fatigue After Stroke (NotFAST) study: results from follow-up six months after stroke. *Top Stroke Rehabil.* 2017;24:592–596. doi: 10.1080/10749357.2017.1368912.
5. Lagogianni C., Thomas S., Lincoln N. Examining the relationship between fatigue and cognition after stroke: a systematic review. *Neuropsychol Rehabil.* 2018;28:57–116. doi: 10.1080/09602011.2015.1127820.
6. Maaijwee N.A., Arntz R.M., Rutten-Jacobs L.C. et al. Post-stroke fatigue and its association with poor functional outcome after stroke in young adults. *J Neurol Neurosurg Psychiatry.* 2015;86:1120–1126. doi: 10.1136/jnnp-2014-308784.
7. Evidence Based Review of Stroke Rehabilitation. Heart and stroke foundation Canadian Partnership for Stroke Recovery. 2018. <http://www.ebrsr.com/> [date access 18.10.2021]

ORCID and contributionship:

Oleksandr Pulyk: 0000-0002-8717-047X^{A-F}

Myroslava Hyryavets: 0000-0001-8419-0590^{A-F}

Taras Studeniak: 0000-0001-6564-1552^{A-F}

Conflict of interest:

The Authors declare no conflict of interest.

CORRESPONDING AUTHOR

Oleksandr Pulyk

Uzhhorod National University

3 Narodna Square, 88000 Uzhhorod, Ukraine

tel: +380506113041

e-mail: apulyk@gmail.com

Received: 17.10.2021

Accepted: 30.04.2022

A – Work concept and design, **B** – Data collection and analysis, **C** – Responsibility for statistical analysis, **D** – Writing the article, **E** – Critical review, **F** – Final approval of the article